

# Université de Nîmes

Laboratoire MIPA  
Université de Nîmes, Site des Carmes  
Place Gabriel Péri, 30021 Nîmes, France  
<http://mipa.unimes.fr>

## On the relaxation of unbounded multiple integrals

by

OMAR ANZA HAFSA AND JEAN-PHILIPPE MANDALLENA

July 2012





# ON THE RELAXATION OF UNBOUNDED MULTIPLE INTEGRALS

OMAR ANZA HAFSA AND JEAN PHILIPPE MANDALLENNA

ABSTRACT. We study the relaxation of multiple integrals of the calculus of variations, where the integrands are nonconvex with convex effective domain and can take the value  $\infty$ . We use local techniques based on measure arguments to prove integral representation in Sobolev spaces of functions which are almost everywhere differentiable. Applications are given in the scalar case and in the case of integrands with quasiconvex growth and  $p(x)$ -growth.

## 1. INTRODUCTION

Let  $m, d \geq 1$  be two integers. Let  $\Omega \subset \mathbb{R}^d$  be a nonempty bounded open set with Lipschitz boundary. Define  $F : W^{1,p}(\Omega; \mathbb{R}^m) \rightarrow [0, \infty]$  by

$$F(u) := \int_{\Omega} f(x, \nabla u(x)) dx,$$

where the integrand  $f : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  is Borel measurable, and  $\mathbb{M}^{m \times d}$  stands for the set of  $m$ -rows and  $d$ -columns matrices. The “relaxed” functional  $\bar{F}$  is given by

$$\bar{F}(u) := \inf \left\{ \liminf_{n \rightarrow \infty} F(u_n) : W^{1,p}(\Omega; \mathbb{R}^m) \ni u_n \rightharpoonup u \right\}$$

(if  $p = \infty$  then replace  $\rightharpoonup$  by  $\overset{*}{\rightharpoonup}$ ). The goal of the paper is to study the integral representation of  $\bar{F}$  for nonconvex integrands  $f$  which can take the value  $\infty$ . In this case, the effective domain  $\text{dom}f(x, \cdot) := \{\xi \in \mathbb{M}^{m \times d} : f(x, \xi) < \infty\}$  of  $f(x, \cdot)$  is the natural set of constraints for the gradients, the interest of such constrained relaxation problems is well described in the book [CDA02].

In the scalar case, i.e., when  $\min\{d, m\} = 1$ , the integral representation of  $\bar{F}$  is studied in [DAMZ04, DAZ05, Zap05]. Under convexity of  $\text{dom}f(x, \cdot)$  and some regularity properties of the multifunction  $x \mapsto \text{dom}f(x, \cdot)$ , integral representations with convexification of  $f(x, \cdot)$  are obtained. The present work focus on the vectorial case, i.e., when  $\min\{d, m\} > 1$ , in this context few is known, in particular the quasiconvexification process when the integrand  $f$  is not finite is not yet understood (for works in this direction see for instance [BB00a, AHM07, AHM08, AH10]). The main difficulty in the integral representation of  $\bar{F}$  is that usually we use an approximation result of functions of  $W^{1,p}(\Omega; \mathbb{R}^m)$  by more regular ones, usually continuous piecewise affine or continuously differentiable functions, and this choice implies different relaxed functionals. This situation is known as Lavrentiev phenomenon (or gap) (see for instance [BB95]). It is not known whether such approximation results exist when no regularity and growth assumptions are made on  $f$  and  $\text{dom}f(x, \cdot)$ . In our work we study the existence of integral representation of  $\bar{F}$  on  $\text{dom}F := \{u \in W^{1,p}(\Omega; \mathbb{R}^m) : F(u) < \infty\}$  without making use of approximation results, and then give some applications showing how to obtain a full integral representation. Following this way, this work is an attempt to establish conditions for

---

*Key words and phrases.* Relaxation, integral representation, constraints.

the existence of integral representation of  $\overline{F}$  with the restrictions that  $\text{dom}f(x, \cdot)$  is convex for all  $x \in \Omega$ ,  $f$  is  $p$ -coercive and  $p \in ]d, \infty]$ . This simplified framework allows us to deal with functions of  $W^{1,p}(\Omega; \mathbb{R}^m)$  that are almost everywhere differentiable in  $\Omega$ , which appears as an important ingredient for the possibility of integral representation of  $\overline{F}$ . The techniques we use are based on measure arguments and localization.

## 2. MAIN RESULTS

We denote by  $\mathcal{O}(\Omega)$  the set of all open subsets of  $\Omega$ . For each  $O \in \mathcal{O}(\Omega)$ , we will denote by  $W_0^{1,p}(O; \mathbb{R}^m)$  the subset of all  $\phi \in W^{1,p}(\Omega; \mathbb{R}^m)$  such that  $\phi = 0$  in  $\Omega \setminus O$  (this definition is equivalent to the classical definition of  $W_0^{1,p}(O; \mathbb{R}^m)$  see [AH96, Chap. 9, p. 233]). We denote by  $Q$  any open cube of  $\mathbb{R}^d$ .

Let  $L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  be a Borel measurable integrand. We consider the following assertions:

(2.1) if  $p \in ]d, \infty[$  then there exists  $c > 0$  such that for every  $(x, \xi) \in \Omega \times \mathbb{M}^{m \times d}$

$$c|\xi|^p \leq L(x, \xi);$$

(2.2) if  $p = \infty$  then there exists  $R_0 > 0$  such that

$$\text{dom}L(x, \cdot) \subset \overline{Q}_{R_0}(0) \text{ a.e. in } \Omega;$$

(2.3) there exists  $\rho_0 > 0$  such that

$$\overline{Q}_{\rho_0}(0) \subset \Lambda_L := \left\{ \xi \in \mathbb{M}^{m \times d} : \int_{\Omega} L(x, \xi) dx < \infty \right\};$$

(2.4) for almost all  $x \in \Omega$

$$\text{dom}L(x, \cdot) \subset \Lambda_L(x) := \left\{ \xi \in \mathbb{M}^{m \times d} : L(x, \xi) = \lim_{\varepsilon \rightarrow 0} \int_{Q_\varepsilon(x)} L(y, \xi) dy \right\};$$

(2.5) there exists  $C > 0$  such that for every  $\xi, \zeta \in \mathbb{M}^{m \times d}$ ,  $x \in \Omega$ , and  $t \in ]0, 1[$

$$L(x, t\xi + (1-t)\zeta) \leq C(1 + L(x, \xi) + L(x, \zeta));$$

(2.6) for almost all  $x \in \Omega$

$$\text{dom}L(x; \cdot) \subset \Xi_L := \left\{ \xi \in \mathbb{M}^{m \times d} : \overline{\lim}_{\delta \rightarrow 0} \omega_\delta^L(\xi) < \infty \right\},$$

$$\text{where } \omega_\delta^L(\xi) := \sup_{\substack{Q \subset \Omega \\ \text{diam}(Q) < \delta}} \inf_{\varphi \in W_0^{1,p}(Q; \mathbb{R}^m)} \int_Q L(x, \xi + \nabla \varphi(x)) dx.$$

*Remark 2.1.* Some remarks on the previous assertions are in order:

(i) The assertions (2.1) and (2.2) are coercivity conditions respectively in the case  $p$  finite and  $p$  non finite, they are used only in the subsection 6.2. Note that if  $p \in ]d, \infty]$  and (2.1) or (2.2) hold, then due to compact embeddings of  $W^{1,p}(\Omega; \mathbb{R}^m)$  in  $L^\infty(\Omega; \mathbb{R}^m)$  we have

$$(2.7) \quad \overline{F}(u) = \inf \left\{ \varliminf_{n \rightarrow \infty} F(u_n) : W^{1,p}(\Omega; \mathbb{R}^m) \ni u_n \xrightarrow{L^\infty} u \right\}.$$

(ii) It is easy to see that the combination ((2.3), (2.4) and (2.6)) is equivalent to: there exists  $\rho_0 > 0$  such that

$$(2.8) \quad \overline{Q}_{\rho_0}(0) \subset \Lambda_L \subset \text{dom}L(x, \cdot) \subset \Lambda_L(x) \cap \Xi_L \quad \text{a.e. in } \Omega.$$

(iii) Due to (2.5), the effective domain  $\text{dom}L(x, \cdot)$  is convex for all  $x \in \Omega$ , the same holds for  $\Lambda_L$  and  $\Lambda_L(x)$ .

(iv) The assertion (2.3) is equivalent to  $0 \in \text{int}(\Lambda_L)$ , where  $\text{int}(\Lambda_L)$  denotes the interior of  $\Lambda_L$ .

We denote by  $Y$  the cube  $] -1, 1[^d$ . Define  $\mathcal{Z}L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  by

$$\mathcal{Z}L(x, \xi) := \liminf_{\varepsilon \rightarrow 0} \inf \left\{ \int_Y L(x + \varepsilon y, \xi + \nabla \varphi(y)) dy : \varphi \in W_0^{1,p}(Y; \mathbb{R}^m) \right\}.$$

*Remark 2.2.* (i) The formula which gives  $\mathcal{Z}L$  can be rewrite

$$(2.9) \quad \mathcal{Z}L(x, \xi) := \liminf_{\varepsilon \rightarrow 0} \inf \left\{ \int_{Q_\varepsilon(x)} L(y, \xi + \nabla \varphi(y)) dy : \varphi \in W_0^{1,p}(Q_\varepsilon(x); \mathbb{R}^m) \right\},$$

where  $Q_\varepsilon(x) = x + \varepsilon Y$  with  $\varepsilon > 0$  and  $x \in \Omega$ .

(ii) If  $L$  does not depend on  $x$ , then  $L$  is  $W^{1,p}$ -quasiconvex in the sense of Ball&Murat[BM84] if and only if  $L = \mathcal{Z}L$ . In fact  $\mathcal{Z}L$  is the generalization to  $x$ -dependent integrand of the Dacorogna quasiconvexification formula. If  $L$  is an Carathéodory integrand with  $p$ -polynomial growth then we can freeze the variable  $x$  and show that

$$\mathcal{Z}L(x, \xi) = \inf \left\{ \int_Y L(x, \xi + \nabla \varphi(y)) dy : \varphi \in W_0^{1,\infty}(Y; \mathbb{R}^m) \right\},$$

which is the Dacorogna quasiconvexification formula for each  $x$  fixed. However the formula (2.9) seems to be the natural generalization when we deal with Borel measurable integrand which can take the value  $\infty$ .

**Definition 2.1.** We say that  $L$  is  $W^{1,p}$ -quasiconvex if  $L = \mathcal{Z}L$ .

We say that  $L$  is radially uniformly upper semicontinuous (ru-usc) if there exists  $a \in L_{\text{loc}}^1(\Omega; ]0, \infty])$  such that  $\lim_{t \rightarrow 1} \overline{\Delta}_L^a(t) \leq 0$ , where  $\Delta_L^a : [0, 1] \rightarrow ]-\infty, \infty]$  is defined by

$$\Delta_L^a(t) := \text{ess sup}_{x \in \Omega} \sup_{\xi \in \text{dom}L(x, \cdot)} \frac{L(x, t\xi) - L(x, \xi)}{a(x) + L(x, \xi)}.$$

The systematic use of the concept of ru-usc functions in the setting of the relaxation of nonconvex functional in the vectorial case start in [AH10], then it was used to prove homogenization results in [AHM11, AHM12].

Define  $\widehat{\mathcal{Z}L} : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  by

$$\widehat{\mathcal{Z}L}(x, \xi) := \lim_{t \rightarrow 1^-} \mathcal{Z}L(x, t\xi).$$

*Remark 2.3.* (i) In fact, if  $L$  is ru-usc then  $\mathcal{Z}L$  too see Lemma 4.7.

(ii) If (2.5) holds and  $\mathcal{Z}L$  is ru-usc then the  $\lim$  in the definition of  $\widehat{\mathcal{Z}L}$  is a limit see Lemma 4.8.

We state the main result of the paper.

**Theorem 2.1.** Assume that  $f$  satisfies (2.1) ((2.2) if  $p = \infty$ ), (2.5) and (2.8). If either  $f$  is ru-usc or  $\mathcal{Z}f$  is ru-usc and  $W^{1,p}$ -quasiconvex, then for every  $u \in \text{dom}F$  we have

$$(2.10) \quad \overline{F}(u) = \int_{\Omega} \widehat{\mathcal{Z}f}(x, \nabla u(x)) dx.$$

*Remark 2.4.* Under the same assumptions the local version of Theorem 2.1 holds also, i.e., if we set

$$\bar{F}(u; O) := \inf \left\{ \liminf_{n \rightarrow \infty} \int_O f(x, \nabla u_n(x)) dx : W^{1,p}(\Omega; \mathbb{R}^m) \ni u_n \rightharpoonup u \right\}$$

then

$$\bar{F}(u; O) = \int_O \widehat{\mathcal{Z}}f(x, \nabla u(x)) dx$$

for all  $O \in \mathcal{O}(\Omega)$  and  $u \in \text{dom}F(\cdot; O)$  where

$$\text{dom}F(\cdot; O) = \left\{ u \in W^{1,p}(\Omega; \mathbb{R}^m) : \int_O f(x, \nabla u(x)) dx < \infty \right\}.$$

(ii) We do not know whether  $\mathcal{Z}f$  is  $W^{1,p}$ -quasiconvex (i.e.,  $\mathcal{Z}(\mathcal{Z}f) = \mathcal{Z}f$ ) when  $f$  is assumed to be ru-usc.

If we consider a stronger assumption (see (2.11)) in place of (2.3) then the following result shows that the full integral representation of  $\bar{F}$  holds.

**Theorem 2.2.** *Assume that  $f$  satisfies (2.1) ((2.2) if  $p = \infty$ ), (2.4), (2.5), (2.6) and there exists  $\rho_0 > 0$  such that for every  $u \in W^{1,p}(\Omega; \mathbb{R}^m)$*

$$(2.11) \quad |u|_{1,p} \leq \rho_0 \implies \int_{\Omega} f(x, \nabla u(x)) dx < \infty.$$

*If either  $f$  is ru-usc or  $\mathcal{Z}f$  is ru-usc and  $W^{1,p}$ -quasiconvex then (2.10) holds for all  $u \in \text{dom}\bar{F}$ .*

*Remark 2.5.* (i) Under the same assumptions the local version of Theorem 2.2 holds also, i.e.,

$$\bar{F}(u; O) = \int_O \widehat{\mathcal{Z}}f(x, \nabla u(x)) dx$$

for all  $O \in \mathcal{O}(\Omega)$  and  $u \in \text{dom}\bar{F}(\cdot; O)$ .

(ii) Theorem 2.2 is mainly used to propose an alternative of the results of [DAMZ04, DAZ05] see subsection 3.2.

(iii) Note that the assertion (2.11) implies (2.3). It seems that condition (2.11) makes sense when  $p = \infty$  because in this case we can show that (2.3) and (2.5) implies (2.11) (see Corollary 4.1). The condition (2.11) means that the effective domain has to be “thick” enough in order that no gap appears when passing from the representation on  $\text{dom}F$  to the representation on  $\text{dom}\bar{F}$ .

Theorems 2.1 and 2.2 are consequences of the following proposition. Define  $\mathcal{Z}F : W^{1,p}(\Omega; \mathbb{R}^m) \times \mathcal{O}(\Omega) \rightarrow [0, \infty]$  by

$$\mathcal{Z}F(u; O) := \inf \left\{ \liminf_{n \rightarrow \infty} \int_O \mathcal{Z}f(x, \nabla u_n(x)) dx : W^{1,p}(\Omega; \mathbb{R}^m) \ni u_n \rightharpoonup u \right\}.$$

**Proposition 2.1.** *Assume that  $f$  satisfies (2.1) ((2.2) if  $p = \infty$ ), (2.3), (2.4) and (2.5). Let  $O \in \mathcal{O}(\Omega)$ .*

(i) *Then for every  $u \in W^{1,p}(\Omega; \mathbb{R}^m)$  such that  $F(tu) < \infty$  for all  $t \in ]0, 1[$ , we have*

$$\bar{F}(tu; O) \leq \int_O \mathcal{Z}f(x, t\nabla u(x)) dx$$

*for all  $t \in ]0, 1[$ .*

(ii) *If  $\mathcal{Z}f$  is ru-usc and  $W^{1,p}$ -quasiconvex then for every  $u \in \text{dom}F$  we have*

$$\bar{\mathcal{Z}}F(u; O) \geq \int_O \widehat{\mathcal{Z}}f(x, \nabla u(x)) dx.$$

(iii) If  $f$  is ru-usc then for every  $u \in \text{dom}F$  we have

$$\overline{F}(u; O) \geq \int_O \widehat{\mathcal{Z}}f(x, \nabla u(x)) dx.$$

In fact the  $\underline{\text{lim}}$  in the definition of  $\mathcal{Z}f$  is a limit if we assume (2.4) and (2.6).

**Proposition 2.2.** *If  $L : \Omega \times \mathbb{M}^{m \times d}$  is a Borel measurable integrand satisfying (2.4) and (2.6) then for almost all  $x \in \Omega$  and every  $\xi \in \text{dom}L(x, \cdot)$*

$$\mathcal{Z}L(x, \xi) = \liminf_{\varepsilon \rightarrow 0} \left\{ \int_Y L(x + \varepsilon y, \xi + \nabla \varphi(y)) dy : \varphi \in W_0^{1,p}(Y; \mathbb{R}^m) \right\}.$$

The plan of paper is as follow. In sect. 3 we give some applications in the case where  $f$  satisfies quasiconvex growth, we show that a full integral representation holds if the functional associated to the quasiconvex growth is sequentially weakly lsc on  $W^{1,p}(\Omega; \mathbb{R}^m)$ . The scalar case is treated by using Theorem 2.2 and adding some assumptions on the regularity of  $\text{dom}f(x, \cdot)$ . Finally an application of Theorem 2.1 is developed in the context of relaxation with integrand satisfying  $p(x)$ -growth.

In sect. 4 we first establish some results on  $L$  and the envelope  $\mathcal{Z}L$  needed for the proof of Proposition 2.1. Then we introduce the concept of ru-usc functionals and state abstract results needed in the proof of Theorem 2.2.

In sect. 5 the proofs of Theorem 2.1 and Theorem 2.2 are given by using Proposition 2.1. The proof of Theorem 2.2 use the abstract result on ru-usc functionals of subsection 4.2 and especially Corollary 4.2.

The sect. 6 and sect. 7 are devoted to the proof of Proposition 2.1. The strategy to prove the upper bound part (i) of Proposition 2.1 is inspired by the paper of [BFM98]. They develop a new method to prove integral representations for relaxed variational functionals and  $\Gamma$ -limit of variational functionals. Roughly, their method consists, when  $\overline{F}(u; \cdot)$  is a Radon measure absolutely continuous with respect to a fixed finite nonnegative Radon measure, to express the Radon-Nikodym derivative of  $\overline{F}(u; \cdot)$  in terms of minima of local Dirichlet problems for  $\overline{F}$ . However, in our case, we use an indirect method for the proof of Proposition 2.1 (i) in the sense that we do not prove directly that  $\overline{F}(u; \cdot)$  is a Radon measure. Note that similar ideas considering the link between the relaxed integrand and minima of local Dirichlet problems appears in [DMM86] and in the context of  $G$ -convergence in [DGS73]. For the proof of the lower bound parts (ii) and (iii) of Proposition 2.1, we use the techniques of localization (also known as blow-up method) and cut-off method introduced by [FM92].

In sect. 8 we give the proof of Proposition 2.2 by using some measure theoretic arguments. More precisely, the proof consists to see  $\mathcal{Z}L(\cdot, \xi)$  as derivate of a set function (see for instance [HK60, Bon72]).

### 3. APPLICATIONS

**3.1. Relaxation with quasiconvex growth.** Let  $p \in ]d, \infty[$ . Let  $G : \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  be a Borel measurable integrand, which is  $W^{1,p}$ -quasiconvex, i.e., for every  $\xi \in \mathbb{M}^{m \times d}$

$$(3.1) \quad G(\xi) = \mathcal{Z}G(\xi).$$

We consider  $G$ -growth on  $f$ , i.e., there exist  $\alpha > 0, \beta > 0$  such that for every  $(x, \xi) \in \Omega \times \mathbb{M}^{m \times d}$

$$(3.2) \quad \alpha G(\xi) \leq f(x, \xi) \leq \beta(1 + G(\xi)).$$

**Theorem 3.1.** *Assume that  $G$  satisfies (2.1), (2.5), (3.1), (3.2) and  $0 \in \text{int}(\text{dom}G)$ . If either  $f$  is ru-usc or  $\mathcal{Z}f$  is  $W^{1,p}$ -quasiconvex and ru-usc then (2.10) holds for all  $u \in W^{1,p}(\Omega; \mathbb{R}^m)$  such that  $\int_{\Omega} G(\nabla u(x))dx < \infty$ . Moreover (2.10) holds for all  $u \in \text{dom}\bar{F}$  if*

$$W^{1,p}(\Omega; \mathbb{R}^m) \ni u \mapsto \int_{\Omega} G(\nabla u(x))dx$$

*is sequentially weakly lower semicontinuous (swlsc) on  $W^{1,p}(\Omega; \mathbb{R}^m)$ .*

*Proof.* By (3.2), it is easy to see that  $\text{dom}f(x, \cdot) = \text{dom}G = \Lambda_f = \Lambda_f(x) = \Xi_f$  a.e. in  $\Omega$ , so  $f$  satisfies (2.8) since  $0 \in \text{int}(\text{dom}G)$ . We have also that  $f$  satisfies (2.1) since  $G$  satisfies (2.1). By Lemma 4.4,  $G$  satisfies (2.5) if and only if  $f$  satisfies (2.5). Applying Theorem 2.1 we obtain (2.10) for all  $u \in \text{dom}F$ . But  $\text{dom}F = \{u \in W^{1,p}(\Omega; \mathbb{R}^m) : \int_{\Omega} G(\nabla u)dx < \infty\}$  since (3.2). If we assume that  $W^{1,p}(\Omega; \mathbb{R}^m) \ni u \mapsto \int_{\Omega} G(\nabla u(x))dx$  is swlsc on  $W^{1,p}(\Omega; \mathbb{R}^m)$ , then again by using (3.2),  $\text{dom}\bar{F} = \{u \in W^{1,p}(\Omega; \mathbb{R}^m) : \int_{\Omega} G(\nabla u)dx < \infty\}$ , thus  $\text{dom}F = \text{dom}\bar{F}$  and the integral representation holds for all  $u \in \text{dom}\bar{F}$ . ■

**3.2. Relaxation in the scalar case.** Let  $L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  be an integrand. We denote by  $L^{**} : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  the convex lower semicontinuous envelope of  $L(x, \cdot)$  for each  $x \in \Omega$ , i.e.,

$$L^{**}(x, \xi) := \sup \{g(x, \xi) : g(x, \cdot) \text{ is convex and lsc, } g(x, \cdot) \leq L(x, \cdot)\}$$

for all  $(x, \xi) \in \Omega \times \mathbb{M}^{m \times d}$ .

To show that the relaxed integrand  $\widehat{\mathcal{Z}L}$  coincides with  $L^{**}$  when  $m = 1$  we need assumption “à la De Arcangelis and all” (see [DAMZ04, DAZ05]). For each  $\varepsilon > 0$  define the multifunction  $D_{\varepsilon} : \Omega \rightrightarrows \mathbb{M}^{m \times d}$  by

$$D_{\varepsilon}(x) := \bigcup_{\varphi \in W_0^{1,p}(\mathbb{Q}_{\varepsilon}(x); \mathbb{R}^m)} \bigcup_{\substack{N \subset \mathbb{Q}_{\varepsilon}(x) \\ |N|=0}} \bigcap_{y \in \mathbb{Q}_{\varepsilon}(x) \setminus N} \text{int}(\text{dom}L(y, \cdot)) - \{\nabla \varphi(y)\},$$

Consider the following assertion:

(3.3) for almost all  $x \in \Omega$

$$\bigcup_{\delta > 0} \bigcap_{\varepsilon \in ]0, \delta[} D_{\varepsilon}(x) \subset \text{dom}L(x, \cdot).$$

**Lemma 3.1.** *If  $L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  is a Borel measurable satisfying (3.3), (2.3), (2.4) and (2.6) then for a.a.  $x \in \Omega$  and for every  $t \in ]0, 1[$  we have*

$$t \text{dom}\widehat{\mathcal{Z}L}(x, \cdot) \subset \text{dom}L(x, \cdot).$$

*Proof.* Fix  $x \in \Omega$  such that (3.3) holds and  $0 \in \text{int}(\text{dom}L(x, \cdot))$ . Fix  $\xi \in \text{dom}\mathcal{Z}L(x, \cdot)$ . Then by Proposition 2.2 there exists  $\varepsilon_0 > 0$  such that

$$\sup_{\varepsilon \in ]0, \varepsilon_0[} \int_{\mathbb{Q}_{\varepsilon}(x)} L(y, \xi + \nabla \varphi_{\varepsilon}(y)) dy < \infty,$$

for some  $\{\varphi_{\varepsilon}\}_{\varepsilon \in ]0, \varepsilon_0[}$ ,  $\varphi_{\varepsilon} \in W_0^{1,p}(\mathbb{Q}_{\varepsilon}(x); \mathbb{R}^m)$ . It follows that for every  $\varepsilon \in ]0, \varepsilon_0[$  there exists a negligible set  $N_x^{\varepsilon} \subset \mathbb{Q}_{\varepsilon}(x)$  such that for every  $y \in \mathbb{Q}_{\varepsilon}(x) \setminus N_x^{\varepsilon}$  we have  $\xi + \nabla \varphi_{\varepsilon}(y) \in \text{dom}L(y, \cdot)$ . It holds  $t\xi + t\nabla \varphi_{\varepsilon}(y) \in t \text{dom}L(y, \cdot)$  for all  $y \in \mathbb{Q}_{\varepsilon}(x) \setminus N_x^{\varepsilon}$  and all  $t \in ]0, 1[$ . Hence  $t\xi \in \bigcap_{y \in \mathbb{Q}_{\varepsilon}(x) \setminus N_x^{\varepsilon}} t \text{dom}L(y, \cdot) - \{t\nabla \varphi_{\varepsilon}(y)\}$  for all  $t \in ]0, 1[$ . By convexity of  $\text{dom}L(y, \cdot)$  and the fact that by (2.3) we have  $0 \in \text{int}(\text{dom}L(y, \cdot))$  for all  $y \in \mathbb{Q}_{\varepsilon}(x) \setminus N'$  for some negligible set  $N'$ , we deduce  $t \text{dom}L(y, \cdot) \subset \text{int}(\text{dom}L(y, \cdot))$  for all  $y \in \mathbb{Q}_{\varepsilon}(x) \setminus N'$  for all  $t \in ]0, 1[$ . It follows that for every  $t \in ]0, 1[$

$$t\xi \in \bigcap_{y \in \mathbb{Q}_{\varepsilon}(x) \setminus (N_x^{\varepsilon} \cup N')} \text{int}(\text{dom}L(y, \cdot)) - \{t\nabla \varphi_{\varepsilon}(y)\}.$$



From (3.3) we deduce that  $t\xi \in \text{dom}L(x, \cdot)$  for all  $t \in ]0, 1[$  which completes the proof.  $\blacksquare$

**Lemma 3.2.** *If the assumptions of Theorem 2.1 and (3.3) hold then for a.a.  $x \in \Omega$  the integrand  $\widehat{\mathcal{Z}L}(x, \cdot)$  is rank-one convex and  $\widehat{\mathcal{Z}L}(x, \cdot) = \overline{\mathcal{Z}L}(x, \cdot)$ , where  $\overline{\mathcal{Z}L}(x, \cdot)$  denotes the lsc envelope of  $\mathcal{Z}L(x, \cdot)$ .*

*Proof.* We have to show that for a.a.  $x \in \Omega$ , for every  $\xi, \zeta \in \mathbb{M}^{m \times d}$  such that  $\text{rank}(\xi - \zeta) \leq 1$  and for every  $\tau \in ]0, 1[$  we have

$$\widehat{\mathcal{Z}L}(x, \tau\xi + (1 - \tau)\zeta) \leq \tau\widehat{\mathcal{Z}L}(x, \xi) + (1 - \tau)\widehat{\mathcal{Z}L}(x, \zeta).$$

Fix  $x_0 \in \Omega'$  where

$$\Omega' := \left\{ x \in \Omega : \forall t \in ]0, 1[ \quad t \text{dom}\widehat{\mathcal{Z}L}(x, \cdot) \subset \text{dom}L(x, \cdot) \subset \Lambda_L(x) \right\}.$$

Since Lemma 3.1 and (2.4) we have  $|\Omega \setminus \Omega'| = 0$ .

Fix  $\xi, \zeta \in \text{dom}\widehat{\mathcal{Z}L}(x_0, \cdot)$ . Thus  $L(x_0, t\xi) < \infty$  and  $L(x_0, t\zeta) < \infty$  for all  $t \in ]0, 1[$ . Fix  $t \in ]0, 1[$ . If  $\chi \in \{\xi, \zeta\}$  then

$$\infty > L(x_0, t\chi) = \lim_{\varepsilon \rightarrow 0} \int_{Q_\varepsilon(x_0)} L(y, t\chi) dy = \lim_{\varepsilon \rightarrow 0} \frac{F(t\chi; Q_\varepsilon(x_0))}{\varepsilon^d}$$

Choose  $\delta_0^t > 0$  such that  $F(t\chi; Q_{\delta_0^t}(x_0)) < \infty$ . By the local version of Theorem 2.1 (with  $O = Q_\delta(x_0)$ ) we have for every  $\varepsilon \in ]0, \delta_0^t[$

$$\overline{F}(t\chi; Q_\varepsilon(x_0)) = \int_{Q_\varepsilon(x_0)} \widehat{\mathcal{Z}L}(y, t\chi) dy.$$

Reasoning as in the proof of the zig-zag lemma (see for instance [BD98, p. 79-80]), we obtain for every  $\tau \in ]0, 1[$

$$\widehat{\mathcal{Z}L}(x_0, \tau t\xi + (1 - \tau)t\zeta) \leq \tau\widehat{\mathcal{Z}L}(x_0, t\xi) + (1 - \tau)\widehat{\mathcal{Z}L}(x_0, t\zeta).$$

The integrand  $\widehat{\mathcal{Z}L}$  is ru-usc

Letting  $t \rightarrow 1$  and using Lemma 4.8 we obtain that  $\widehat{\mathcal{Z}L}(x_0, \cdot)$  is rank-one convex. Then  $\widehat{\mathcal{Z}L}(x_0, \cdot)$  is separately convex and so is continuous in  $\text{int}(\text{dom}\widehat{\mathcal{Z}L}(x_0, \cdot))$  see [Dac08, Theorem 2.31, p. 47]. Applying Lemma 4.7 together with Theorem 4.2 the function  $\widehat{\mathcal{Z}L}(x_0, \cdot)$  is the lsc envelope of  $\mathcal{Z}L(x_0, \cdot)$ .  $\blacksquare$

The integral representation of  $\overline{F}$  was studied in [DAMZ04, DAZ05] in the scalar case with  $p = \infty$ , we propose here the following alternative result.

**Theorem 3.2.** *Assume that  $m = 1$ . Assume that  $f$  satisfies (2.1) ((2.2) if  $p = \infty$ ), (2.4), (2.5), (2.6), (2.11) and (3.3). If either  $f$  is ru-usc or  $\mathcal{Z}f$  is  $W^{1,p}$ -quasiconvex and ru-usc then for every  $u \in \text{dom}\overline{F}$  we have*

$$\overline{F}(u) = \int_{\Omega} f^{**}(x, \nabla u(x)) dx.$$

Moreover  $f^{**}(x, \cdot) = \overline{\mathcal{Z}f}(x, \cdot) = \widehat{\mathcal{Z}f}(x, \cdot)$  a.e. in  $\Omega$ .

*Proof.* Applying Theorem 2.2 the representation (2.10) holds for all  $u \in \text{dom}\overline{F}$ . Fix  $\xi \in \mathbb{M}^{1 \times d}$ . On one hand, by a well known lower semicontinuity result (see for instance [But89, Theorem 4.1.1]) we have for every  $O \in \mathcal{O}(\Omega)$  and  $u \in W^{1,p}(\Omega)$

$$\begin{aligned} \overline{F}(u; O) &\geq \inf \left\{ \liminf_{n \rightarrow \infty} \int_O f^{**}(x, \nabla u_n(x)) dx : W^{1,p}(\Omega) \ni u_n \rightharpoonup u \right\} \\ &\geq \int_O f^{**}(x, \nabla u(x)) dx. \end{aligned}$$

It follows that  $\widehat{\mathcal{Z}}f(x, \cdot) \geq f^{**}(x, \cdot)$  a.e. in  $\Omega$ . On the other hand by Lemma 3.2 we have  $\overline{\mathcal{Z}}f(x, \cdot)$  is convex and lsc and  $f(x, \cdot) \geq \overline{f}(x, \cdot) \geq \overline{\mathcal{Z}}f(x, \cdot) = \widehat{\mathcal{Z}}f(x, \cdot) \geq f^{**}(x, \cdot)$  a.e.  $\Omega$ , and the proof is complete.  $\blacksquare$

*Remark 3.1.* If we consider the case  $p = \infty$  in Theorem 3.2, we can replace the assumption (2.11) by (2.3) since (2.3) and (2.5) implies (2.11) see Corollary 4.1.

**3.3. Relaxation with  $p(x)$ -growth.** Let  $p \in ]d, \infty[$ . Let  $p : \Omega \rightarrow [0, \infty[$  be a measurable function such that  $p \leq p(x)$  for all  $x \in \Omega$ .

Let  $f : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty[$  be a Borel measurable integrand.

Consider the assertions:

(3.4) for each  $\xi \in \mathbb{M}^{m \times d}$  we have

$$|\xi|^{p(\cdot)} \in L^1(\Omega) \quad \text{and} \quad \overline{\lim}_{\delta \rightarrow 0} \sup_{\substack{Q \subset \Omega, \text{ cube} \\ \text{diam}(Q) < \delta}} \int_Q |\xi|^{p(x)} dx < \infty;$$

(3.5) there exist  $\alpha, \beta > 0$  such that for every  $(x, \xi) \in \Omega \times \mathbb{M}^{m \times d}$  we have

$$\alpha |\xi|^{p(x)} \leq f(x, \xi) \leq \beta(1 + |\xi|^{p(x)}).$$

When (3.5) holds, we say that  $f$  has  $p(x)$ -growth. The condition (3.4) is satisfied if  $p(\cdot) \leq p^*$  for some  $p^* \in ]d, \infty[$ .

**Theorem 3.3.** *Assume that (3.5) holds. If (2.10) holds for each  $u \in \text{dom} \overline{F}$ , then  $\widehat{\mathcal{Z}}f$  is a Carathéodory integrand which is ru-usc and rank-one convex with respect to the second variable.*

*Proof.* Reasoning as in the proof of the zig-zag lemma (see for instance [BD98, p. 79-80]), we obtain that  $\widehat{\mathcal{Z}}f(x, \cdot)$  is rank-one convex for all  $x \in \Omega$ . Then  $\widehat{\mathcal{Z}}f(x, \cdot)$  is separately convex, moreover using (3.4), it is to see that  $\widehat{\mathcal{Z}}f$  satisfies: for a.a.  $x \in \Omega$  and for every  $\xi \in \mathbb{M}^{m \times d}$  it holds

$$(3.6) \quad \alpha |\xi|^{p(x)} \leq \widehat{\mathcal{Z}}f(x, \xi) \leq \beta(1 + |\xi|^{p(x)}).$$

So by using [Dac08, Theorem 2.31, p. 47] we obtain that for a.a.  $x \in \Omega$  the function  $\widehat{\mathcal{Z}}f(x, \cdot)$  is continuous in  $\text{int}(\text{dom} \widehat{\mathcal{Z}}L(x, \cdot))$ . But for a.a.  $x \in \Omega$  we have  $\text{int}(\text{dom} \widehat{\mathcal{Z}}f(x, \cdot)) = \text{dom} \widehat{\mathcal{Z}}f(x, \cdot) = \mathbb{M}^{m \times d}$  since (3.6).

From (3.6) and [Dac08, Prop. 2.32, p. 51] there exists  $K > 0$  such that for a.a.  $x \in \Omega$ , every  $t \in ]0, 1[$  and every  $\xi \in \mathbb{M}^{m \times d}$

$$\begin{aligned} |\widehat{\mathcal{Z}}f(x, t\xi) - \widehat{\mathcal{Z}}f(x, \xi)| &\leq K|t\xi - \xi| \left( 1 + |t\xi|^{p(x)-1} + |\xi|^{p(x)-1} \right) \\ &\leq (1-t)4K \left( 1 + |\xi|^{p(x)} \right) \\ &\leq (1-t)4K \left( 1 + \frac{1}{\alpha} \widehat{\mathcal{Z}}f(x, \xi) \right) \\ &\leq (1-t)4K \left( 1 + \frac{1}{\alpha} \right) \left( 1 + \widehat{\mathcal{Z}}f(x, \xi) \right), \end{aligned}$$

where we used (3.5). We obtain  $\Delta_{\widehat{\mathcal{Z}}f}^1(t) \leq (1-t)4K(1 + \frac{1}{\alpha})$  which shows that  $\widehat{\mathcal{Z}}f$  is ru-usc by letting  $t \rightarrow 1$ .  $\blacksquare$

**Theorem 3.4.** *Assume that (3.4) and (3.5) hold. If  $\mathcal{Z}f$  is  $W^{1,p}$ -quasiconvex and ru-usc, then (2.10) holds for each  $u \in \text{dom} \overline{F}$ .*

*Proof.* Since (3.5) and (3.4) we have  $\text{dom}f(x, \cdot) = \mathbb{M}^{m \times d}$  for all  $x \in \Omega$  and (2.8) holds. Moreover, (2.1) holds since  $p(\cdot) \geq p > d$ . It is easy to check that (2.5) holds since Lemma 4.4 and the fact that for each  $x$  the function  $\xi \mapsto |\xi|^{p(x)}$  is convex. Apply Theorem 2.1 we have (2.10) for all  $u \in \text{dom}F$ . Using convexity it is easy to see that the functional  $W^{1,p}(\Omega; \mathbb{R}^m) \ni u \mapsto \int_{\Omega} |\nabla u(x)|^{p(x)} dx$  is swlsc on  $W^{1,p}(\Omega; \mathbb{R}^m)$ . Hence

$$\text{dom}\bar{F} = \text{dom}F = \left\{ u \in W^{1,p}(\Omega; \mathbb{R}^m) : \int_{\Omega} |\nabla u(x)|^{p(x)} dx < \infty \right\},$$

and the proof is complete.  $\blacksquare$

#### 4. PRELIMINARY RESULTS

**4.1. Some properties of  $L$  and  $\mathcal{Z}L$ .** The following lemma is an extension for nonconvex functions satisfying (2.3) and (2.5) of the classical local upper bound property for convex functions.

**Lemma 4.1.** *Let  $L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  be a Borel measurable integrand. If  $L$  satisfies (2.3) and (2.5) then*

$$\int_{\Omega} \mathcal{M}_0(x) dx < \infty \quad \text{where} \quad \mathcal{M}_0(\cdot) := \sup_{\zeta \in \bar{Q}_{\rho_0}(0)} L(\cdot, \zeta).$$

*Proof.* Each matrix  $\xi \in \bar{Q}_{\rho_0}(0)$  is identified to the vector

$$\xi = (\xi_{11}, \dots, \xi_{1d}, \dots, \xi_{i1}, \dots, \xi_{id}, \dots, \xi_{m1}, \dots, \xi_{md}).$$

Consider the finite subset  $\mathcal{S} := \{(\xi_{11}, \dots, \xi_{md}) : \xi_{ij} \in \{-\rho_0, 0, \rho_0\}\} \subset \bar{Q}_{\rho_0}(0)$  and define the function  $L^* : \Omega \rightarrow [0, \infty]$  by  $L^*(x) := \max_{\xi \in \mathcal{S}} L(x, \xi)$ . The function  $L^*$  belongs to  $L^1(\Omega)$  since (2.3), indeed, for each  $x \in \Omega$  choose one  $\xi_x \in \mathcal{S}$  such that  $L^*(x) = L(x, \xi_x)$ , and for each  $\xi \in \mathcal{S}$  consider the sets  $M_{\xi} := \{y \in \Omega : \xi_y = \xi\}$ , then the finite family  $\{M_{\xi}\}_{\xi \in \mathcal{S}}$  is pairwise disjoint,  $\Omega = \cup_{\xi \in \mathcal{S}} M_{\xi}$  and

$$\int_{\Omega} L^*(x) dx = \sum_{\xi \in \mathcal{S}} \int_{M_{\xi}} L(x, \xi_x) dx \leq \sum_{\xi \in \mathcal{S}} \int_{\Omega} L(x, \xi) dx < \infty.$$

Fix  $x \in \Omega$ . Let  $\zeta = (\zeta_{11}, \dots, \zeta_{1d}, \dots, \zeta_{i1}, \dots, \zeta_{id}, \dots, \zeta_{m1}, \dots, \zeta_{md}) \in \mathcal{S}$  with  $\xi_{ij} = \zeta_{ij}$  for all  $i \neq 1$  and  $j \neq 1$ . If  $\xi_{ij} = \zeta_{ij}$  for all  $i \neq 1$  and  $j \neq 1$ , and  $\xi_{11} \neq 0$ , then by (2.5) we have

$$\begin{aligned} (4.1) \quad L(x, \xi) &= L\left(x, \frac{|\xi_{11}|}{\rho_0} \text{sgn}(\xi_{11}) \rho_0 + \left(1 - \frac{|\xi_{11}|}{\rho_0}\right) 0, \dots, \xi_{1d}, \dots, \xi_{m1}, \dots, \xi_{md}\right) \\ &\leq C(1 + L(x, \rho_0, \dots, \xi_{md}) + L(x, 0, \dots, \xi_{md})) \\ &\leq 2C(1 + L^*(x)), \end{aligned}$$

where  $\text{sgn}(\xi_{ij})$  denotes the sign of  $\xi_{ij}$ . The same upper bound in (4.1) holds for  $L(x, \xi)$  when  $\xi_{11} = 0$ .

Assume now that  $\xi_{ij} = \zeta_{ij}$  for all  $i \neq 1$  and  $j \notin \{1, 2\}$ . Then by using (4.1) and (2.5), we have

$$\begin{aligned} L(x, \xi) &= L\left(x, \xi_{11}, \frac{|\xi_{12}|}{\rho_0} \text{sgn}(\xi_{12}) \rho_0 + \left(1 - \frac{|\xi_{12}|}{\rho_0}\right) 0, \dots, \xi_{1d}, \dots, \xi_{m1}, \dots, \xi_{md}\right) \\ &\leq C(1 + 2C(1 + L^*(x)) + L^*(x)). \\ &\leq C(1 + 2C)(1 + L^*(x)). \end{aligned}$$

Recursively, we obtain  $C^* > 0$  which depends on  $C$  only, such that

$$L(x, \xi) \leq C^*(1 + L^*(x))$$

for all  $(x, \xi) \in \Omega \times \overline{Q}_{\rho_0}(0)$ . Integrating over  $\Omega$  we obtain the result

$$\int_{\Omega} \sup_{\zeta \in \overline{Q}_{\rho_0}(0)} L(x, \zeta) dx \leq C^* \left( |\Omega| + \int_{\Omega} L^*(x) dx \right) < \infty.$$

■

**Corollary 4.1.** *If  $p = \infty$  then (2.3) and (2.5) implies (2.11), i.e., there exists  $\rho_0 > 0$  such that  $\int_{\Omega} L(x, \nabla u(x)) dx < \infty$  whenever  $|u|_{1, \infty} \leq \rho_0$  for all  $u \in W^{1, \infty}(\Omega; \mathbb{R}^m)$ .*

*Proof.* It is enough to remark that  $\nabla u(\cdot) \in \overline{Q}_{\rho_0}(0)$  a.e. in  $\Omega$  whenever  $|u|_{1, \infty} \leq \rho_0$ , and then to apply Lemma 4.1. ■

For the proof of Theorem 2.1 we need the following result.

**Lemma 4.2.** *If  $L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  is a Borel measurable integrand which satisfies (2.4) then for a.a.  $x \in \Omega$  and every  $\xi \in \mathbb{M}^{m \times d}$  we have  $\mathcal{Z}L(x, \xi) \leq L(x, \xi)$ .*

*Proof.* Fix  $x_0 \in \Omega'$  where  $\Omega' := \{x \in \Omega : \text{dom}L(x, \cdot) \subset \Lambda_L(x)\}$ . We have  $|\Omega \setminus \Omega'| = 0$  since (2.4). If  $\xi \notin \text{dom}L(x_0, \cdot)$  then  $\mathcal{Z}L(x_0, \xi) \leq \infty = L(x_0, \xi)$ . Now, if  $\xi \in \text{dom}L(x_0, \cdot)$  then by (2.4)  $\lim_{\varepsilon \rightarrow 0} \int_{Q_{\varepsilon}(x_0)} L(z, \xi) dz = L(x_0, \xi)$ . Using the definition of  $\mathcal{Z}L$  we finish the proof. ■

**Lemma 4.3.** *If  $L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  is a Borel measurable integrand which satisfies (2.5) then  $\mathcal{Z}L$  satisfies (2.5).*

*Proof.* Let  $x \in \Omega$ ,  $\xi, \zeta \in \mathbb{M}^{m \times d}$  and  $t \in ]0, 1[$ . There exist  $\{\varphi_{\xi, \varepsilon}\}_{\varepsilon}, \{\varphi_{\zeta, \varepsilon}\}_{\varepsilon} \subset W_0^{1, p}(Q_{\varepsilon}(x); \mathbb{R}^m)$  such that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_{Q_{\varepsilon}(x)} L(y, \xi + \nabla \varphi_{\xi, \varepsilon}) dy &= \mathcal{Z}L(x, \xi) \\ \lim_{\varepsilon \rightarrow 0} \int_{Q_{\varepsilon}(x)} L(y, \zeta + \nabla \varphi_{\zeta, \varepsilon}) dy &= \mathcal{Z}L(x, \zeta). \end{aligned}$$

Since  $\varphi_{\varepsilon} := t\varphi_{\xi, \varepsilon} + (1-t)\varphi_{\zeta, \varepsilon} \in W_0^{1, p}(Q_{\varepsilon}(x); \mathbb{R}^m)$  we have

$$\begin{aligned} \mathcal{Z}L(x, t\xi + (1-t)\zeta) &\leq \liminf_{\varepsilon \rightarrow 0} \int_{Q_{\varepsilon}(x)} L(y, t\xi + (1-t)\zeta + \nabla \varphi_{\varepsilon}) dy \\ &\leq C \liminf_{\varepsilon \rightarrow 0} \int_{Q_{\varepsilon}(x)} (1 + L(y, \xi + \nabla \varphi_{\xi, \varepsilon}) + L(y, \zeta + \nabla \varphi_{\zeta, \varepsilon})) dy \\ &\leq C(1 + \mathcal{Z}L(x, \xi) + \mathcal{Z}L(x, \zeta)), \end{aligned}$$

which completes the proof. ■

The following result shows that the condition (2.5) is shared by integrands with the same growth.

**Lemma 4.4.** *If  $L_1, L_2 : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  are two Borel measurable integrands such that for some  $\alpha, \beta > 0$  and for every  $(x, \xi) \in \Omega \times \mathbb{M}^{m \times d}$  it holds*

$$\alpha L_2(x, \xi) \leq L_1(x, \xi) \leq \beta(1 + L_2(x, \xi)),$$

*then  $L_1$  satisfies (2.5) if and only if  $L_2$  satisfies (2.5).*

*Proof.* Assume that  $L_1$  satisfies (2.5). Fix  $x \in \Omega$ . Let  $\xi, \zeta \in \mathbb{M}^{m \times d}$  and  $t \in ]0, 1[$ . Then

$$\begin{aligned} L_2(x, t\xi + (1-t)\zeta) &\leq \frac{1}{\alpha} L_1(x, t\xi + (1-t)\zeta) \\ &\leq \frac{C}{\alpha} (1 + L_1(x, \xi) + L_1(x, \zeta)) \\ &\leq \frac{C}{\alpha} (1 + 2\beta + \beta L_2(x, \xi) + \beta L_2(x, \zeta)) \\ &\leq \frac{C}{\alpha} (1 + 2\beta) (1 + L_2(x, \xi) + L_2(x, \zeta)). \end{aligned}$$

In the same manner we can verify that if  $L_2$  satisfies (2.5) then  $L_1$  too.  $\blacksquare$

**4.2. Ru-usc functionals.** Let  $(X, \tau)$  be a topological vector space and  $J : X \rightarrow [0, \infty]$  be a function. For each  $a > 0$  and  $D \subset \text{dom} J$  we define  $\Delta_{J,D}^a : [0, 1] \rightarrow ]-\infty, \infty]$  by

$$\Delta_{J,D}^a(t) := \sup_{u \in D} \frac{J(tu) - J(u)}{a + J(u)}.$$

When  $D = \text{dom} J$  we will write  $\Delta_J^a := \Delta_{J,D}^a$ .

**Definition 4.1.** Given  $D \subset \text{dom} J$ , we say that  $J$  is ru-usc in  $D$ , if there exists  $a > 0$  such that

$$\overline{\lim}_{t \rightarrow 1} \Delta_{J,D}^a(t) \leq 0.$$

*Remark 4.1.* If  $J$  is ru-usc in  $D$  then

$$(4.2) \quad \overline{\lim}_{t \rightarrow 1} J(tu) \leq J(u)$$

for all  $u \in D$ . Indeed, given  $u \in D$ , we have

$$J(tu) \leq \Delta_{J,D}^a(t) (a + J(u)) + J(u) \text{ for all } t \in [0, 1],$$

which gives (4.2) since  $a + J(u) > 0$  and  $\overline{\lim}_{t \rightarrow 1} \Delta_J^a(t) \leq 0$ .

*Remark 4.2.* If there exists  $u_0 \in D$  such that  $J$  is ‘‘radially’’ lower semicontinuous at  $u_0$  in the sense that

$$(4.3) \quad \underline{\lim}_{t \rightarrow 1} J(tu_0) - J(u_0) \geq 0.$$

Then

$$(4.4) \quad \underline{\lim}_{t \rightarrow 1} \Delta_{J,D}^a(t) \geq 0$$

for all  $a > 0$ . Indeed, given such  $u \in D$ , for any  $a > 0$  we have

$$\Delta_{J,D}^a(t) \geq \frac{J(tu_0) - J(u_0)}{a + J(u_0)} \text{ for all } t \in [0, 1],$$

which gives (4.4) since  $a + J(u_0) > 0$  and (4.3).

For a subset  $D \subset X$ , we denote by  $\overline{D}^\tau$  the closure of  $D$  with respect to  $\tau$ .

**Lemma 4.5.** Let  $D \subset \text{dom} J$  be a  $\tau$ -star shaped subset with respect 0, i.e.,

$$(4.5) \quad t\overline{D}^\tau \subset D \text{ for all } t \in ]0, 1[.$$

If  $J$  is ru-usc in  $D$  then

$$\underline{\lim}_{t \rightarrow 1} J(tu) = \overline{\lim}_{t \rightarrow 1} J(tu)$$

for all  $u \in \overline{D}^\tau$ .

*Proof.* Fix  $u \in \overline{D}^\tau$ . It suffices to prove that

$$(4.6) \quad \overline{\lim}_{t \rightarrow 1} J(tu) \leq \underline{\lim}_{t \rightarrow 1} J(tu).$$

Without loss of generality we can assume that  $\underline{\lim}_{t \rightarrow 1} J(tu) < \infty$  and there exist  $\{t_n\}_n, \{s_n\}_n \subset ]0, 1[$  such that:

- $t_n \rightarrow 1, s_n \rightarrow 1$  and  $\frac{t_n}{s_n} \rightarrow 1$ ;
- $\overline{\lim}_{t \rightarrow 1} J(tu) = \lim_{n \rightarrow \infty} J(t_n u)$ ;
- $\underline{\lim}_{t \rightarrow 1} J(tu) = \lim_{n \rightarrow \infty} J(s_n u)$ .

From (4.5) we see that for every  $n \geq 1$ ,  $s_n u \in D$ , and so we can assert that for every  $n \geq 1$ ,

$$(4.7) \quad J(t_n u) \leq a \Delta_{J,D}^a \left( \frac{t_n}{s_n} \right) + \left( 1 + \Delta_{J,D}^a \left( \frac{t_n}{s_n} \right) \right) J(s_n u).$$

On the other hand, as  $J$  is ru-usc in  $D$  we have  $\overline{\lim}_{n \rightarrow \infty} \left( 1 + \Delta_{J,D}^a \left( \frac{t_n}{s_n} \right) \right) \leq 1$  and  $\overline{\lim}_{n \rightarrow \infty} a \Delta_{J,D}^a \left( \frac{t_n}{s_n} \right) \leq 0$  since  $a > 0$ , and (4.6) follows from (4.7) by letting  $n \rightarrow \infty$ .  $\blacksquare$

Define  $\widehat{J} : X \rightarrow [0, \infty]$  by

$$\widehat{J}(u) := \underline{\lim}_{t \rightarrow 1} J(tu).$$

**Lemma 4.6.** *If  $J$  is ru-usc in a  $\tau$ -star shaped set  $D \subset \text{dom} J$  then  $\widehat{J}$  is ru-usc in  $\overline{D}^\tau \cap \text{dom} \widehat{J}$ .*

*Proof.* Fix any  $t \in ]0, 1[$  and any  $u \in \overline{D}^\tau \cap \text{dom} \widehat{J}$ . We have  $tu \in D$  since (4.5) holds. From Lemma 4.5 we can assert that:

- $\widehat{J}(u) = \lim_{s \rightarrow 1} J(su)$ ;
- $\widehat{J}(tu) = \lim_{s \rightarrow 1} J(s(tu))$ ,

and consequently

$$(4.8) \quad \frac{\widehat{J}(tu) - \widehat{J}(u)}{a + \widehat{J}(u)} = \lim_{s \rightarrow 1} \frac{J(t(su)) - J(su)}{a + J(su)}.$$

On the other hand, by (4.5) we have  $su \in D$  for all  $s \in ]0, 1[$ , and so

$$\frac{J(t(su)) - J(su)}{a + J(su)} \leq \Delta_{L,D}^a(t) \text{ for all } s \in ]0, 1[.$$

Letting  $s \rightarrow 1$  and using (4.8) we deduce that  $\Delta_{\widehat{J}, \overline{D}^\tau \cap \text{dom} \widehat{J}}^a(t) \leq \Delta_{J,D}^a(t)$  for all  $t \in ]0, 1[$ , which implies that  $\widehat{J}$  is ru-usc in  $\overline{D}^\tau \cap \text{dom} \widehat{J}$  since  $J$  is ru-usc in  $D$ .  $\blacksquare$

**Theorem 4.1.** *If  $J$  is ru-usc in a  $\tau$ -star shaped set  $D \subset \text{dom} J$ , and  $\tau$  sequentially lower semicontinuous on  $D$  then:*

- (i)  $\widehat{J}(u) = \begin{cases} J(u) & \text{if } u \in D \\ \underline{\lim}_{t \rightarrow 1} J(tu) & \text{if } u \in \overline{D}^\tau \setminus D \end{cases}$
- (ii)  $\widehat{J} = \overline{J}^D$  on  $\overline{D}^\tau$  where

$$\overline{J}^D(u) := \inf \left\{ \underline{\lim}_{n \rightarrow \infty} J(u_n) : D \ni u_n \xrightarrow{\tau} u \right\}$$

*Proof.* (i) Lemma 4.5 shows that, for  $u \in \overline{D}^\tau$ ,  $\widehat{J}(u) = \lim_{t \rightarrow 1} J(tu)$ . From remark 4.1 we see that if  $u \in D$  then  $\overline{\lim}_{t \rightarrow 1} J(tu) \leq J(u)$ . On the other hand, from (4.5) it follows that if  $u \in D$  then  $tu \in D$  for all  $t \in ]0, 1[$ . Thus,  $\underline{\lim}_{t \rightarrow 1} J(tu) \geq J(u)$  whenever  $u \in D$  since  $J$  is  $\tau$  lsc on  $D$ , and (i) follows.

(ii) Let  $u \in \overline{D}^\tau$ . Using (4.5) we have  $tu \in D$  for all  $t \in ]0, 1[$ , so by Lemma 4.1 and lower semicontinuity it follows that  $\widehat{J}(u) = \lim_{t \rightarrow 1} J(tu) \geq \overline{J}^D(u)$ . It remains to prove that

$$(4.9) \quad \overline{J}^D(u) \geq \widehat{J}(u).$$

Choose a sequence  $\{u_n\}_n \subset D$  such that  $u_n \xrightarrow{\tau} u$  and  $\lim_{n \rightarrow \infty} J(u_n) = \overline{J}^D(u)$ . By (4.5) we see that, for any  $t \in ]0, 1[$ ,  $tu_n \in D$  for all  $n \geq 1$ , and consequently

$$\underline{\lim}_{n \rightarrow \infty} J(tu_n) \geq J(tu) \text{ for all } t \in ]0, 1[$$

because  $J$  is  $\tau$  lsc on  $D$ . It follows that

$$(4.10) \quad \overline{\lim}_{t \rightarrow 1} \underline{\lim}_{n \rightarrow \infty} J(tu_n) \geq \widehat{J}(u).$$

On the other hand, for every  $n \geq 1$  and every  $t \in [0, 1]$ , we have

$$J(tu_n) \leq (1 + \Delta_{J,D}^a(t))J(u_n) + a\Delta_{J,D}^a(t).$$

As  $J$  is ru-usc in  $D$ , letting  $n \rightarrow \infty$  and  $t \rightarrow 1$  we obtain

$$\overline{\lim}_{t \rightarrow 1} \underline{\lim}_{n \rightarrow \infty} J(tu_n) \leq \lim_{n \rightarrow \infty} J(u_n) = \overline{J}^D(u),$$

which gives (4.9) when combined with (4.10).  $\blacksquare$

The following result is a consequence of Theorem 4.1 when  $J = \overline{F}$  where  $F : X \rightarrow [0, \infty]$  is a functional, and  $\overline{F} : X \rightarrow [0, \infty]$  is the  $\tau$  sequential lsc envelope defined by

$$\overline{F}(u) := \inf \left\{ \underline{\lim}_{n \rightarrow \infty} F(u_n) : X \ni u_n \xrightarrow{\tau} u \right\}.$$

**Corollary 4.2.** *Assume that  $\text{dom}F$  is  $\tau$ -star shaped with respect to 0,  $\overline{F}$  is ru-usc in  $\text{dom}F$ , and  $\overline{F} = I$  on  $\text{dom}F$  where  $I : X \rightarrow [0, \infty]$  is a functional. Then*

$$(4.11) \quad \overline{F}(u) := \begin{cases} I(u) & \text{if } u \in \text{dom}F \\ \lim_{t \rightarrow 1} I(tu) & \text{if } u \in \text{dom}\overline{F} \setminus \text{dom}F \\ \infty & \text{otherwise.} \end{cases}$$

**4.2.1. Ru-usc integrands.** Let  $U \subset \mathbb{R}^d$  be an measurable set and let  $L : M \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  be a measurable integrand. For each  $x \in M$  and for each  $a \in L_{\text{loc}}^1(U; ]0, \infty])$ , we define  $\Delta_L^a : [0, 1] \rightarrow ]-\infty, \infty]$  by

$$\Delta_L^a(t) := \text{ess sup}_{x \in U} \sup_{\xi \in \text{dom}L(x, \cdot)} \frac{L(x, t\xi) - L(x, \xi)}{a(x) + L(x, \xi)}.$$

**Definition 4.2.** *We say that  $L$  is radially uniformly upper semicontinuous (ru-usc) if there exists  $a \in L_{\text{loc}}^1(U; ]0, \infty])$  such that*

$$\overline{\lim}_{t \rightarrow 1} \Delta_L^a(t) \leq 0.$$

*Remark 4.3.* If  $L$  is ru-usc then

$$(4.12) \quad \overline{\lim}_{t \rightarrow 1} L(x, t\xi) \leq L(x, \xi)$$

for all  $x \in U$  and all  $\xi \in \text{dom}L(x, \cdot)$ .

*Remark 4.4.* If there exist  $x \in U$  and  $\xi \in \text{dom}L(x, \cdot)$  such that  $L(x, \cdot)$  is lsc at  $\xi$  then

$$(4.13) \quad \varliminf_{t \rightarrow 1} \Delta_L^a(t) \geq 0$$

for all  $a \in L_{\text{loc}}^1(U; ]0, \infty])$ .

**Lemma 4.7.** *If  $L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  is a ru-usc Borel measurable integrand then  $ZL$  is ru-usc and  $\Delta_{ZL}^a(t) \leq \Delta_L^a(t)$  for all  $t \in ]0, 1[$ .*

*Proof.* Indeed, fix  $x \in \Omega$  such that it is a Lebesgue point for the function  $a \in L^1(\Omega)$  which appears in the definition of the ru-usc for  $L$ . Fix  $\xi \in \text{dom}ZL(x, \cdot)$  and choose a sequence  $\{\varphi_\varepsilon\}_\varepsilon \subset W_0^{1,p}(\mathbb{Q}_\varepsilon(x); \mathbb{R}^m)$  satisfying for every  $\varepsilon > 0$

$$\varepsilon + ZL(x, \xi) \geq \int_{\mathbb{Q}_\varepsilon(x)} L(y, \xi + \nabla \varphi_\varepsilon) dy.$$

Fix  $\varepsilon > 0$ . Then  $\xi + \nabla \varphi_\varepsilon(y) \in \text{dom}L(y, \cdot)$  a.e. in  $\mathbb{Q}_\varepsilon(x)$ . We have for every  $t \in ]0, 1[$

$$\begin{aligned} & \int_{\mathbb{Q}_\varepsilon(x)} L(y, t\xi + \nabla \varphi_\varepsilon) dy - ZL(x, \xi) \\ & \leq \int_{\mathbb{Q}_\varepsilon(x)} L(y, t\xi + \nabla \varphi_\varepsilon) dy - \int_{\mathbb{Q}_\varepsilon(x)} L(y, \xi + \nabla \varphi_\varepsilon) dy + \varepsilon \\ & \leq \Delta_L(t) \left( \int_{\mathbb{Q}_\varepsilon(x)} a(y) + L(y, \xi + \nabla \varphi_\varepsilon) dy \right) + \varepsilon \\ & \leq \Delta_L(t) \left( \int_{\mathbb{Q}_\varepsilon(x)} a(y) dy + \varepsilon + ZL(x, \xi) \right) + \varepsilon \end{aligned}$$

Taking the infimum over all  $\varphi \in W_0^{1,p}(\mathbb{Q}_\varepsilon(x); \mathbb{R}^m)$  and passing to the limit  $\varepsilon \rightarrow 0$  we obtain  $\Delta_{ZL}^a(t) \leq \Delta_L^a(t)$  for all  $t \in ]0, 1[$ , and the proof is complete.  $\blacksquare$

Define  $\widehat{L} : U \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  by

$$\widehat{L}(x, \xi) := \varliminf_{t \rightarrow 1} L(x, t\xi).$$

The proof of the following result is similar to the proof of Lemma 4.5 and Lemma 4.6 (see [AHM11]).

**Lemma 4.8.** *If  $L$  is ru-usc and for a.a.  $x \in U$  the effective domain  $\text{dom}L(x, \cdot)$  is star shaped with respect to 0, i.e.,  $t\overline{\text{dom}L(x, \cdot)} \subset \text{dom}L(x, \cdot)$  for all  $t \in ]0, 1[$  then  $\widehat{L}$  is ru-usc and*

$$\widehat{L}(x, \xi) = \lim_{t \rightarrow 1} L(x, t\xi).$$

We can now state the analogue of Theorem 4.1 (see [AHM11]).

**Theorem 4.2.** *If  $L$  is ru-usc and if for every  $x \in U$ ,*

$$t\overline{\text{dom}L(x, \cdot)} \subset \text{int}(\text{dom}L(x, \cdot)) \text{ for all } t \in ]0, 1[$$

and  $L(x, \cdot)$  is lsc on  $\text{int}(\text{dom}L(x, \cdot))$ , then:

$$(i) \quad \widehat{L}(x, \xi) = \begin{cases} L(x, \xi) & \text{if } \xi \in \text{int}(\text{dom}L(x, \cdot)) \\ \lim_{t \rightarrow 1} L(x, t\xi) & \text{if } \xi \in \partial \text{dom}L(x, \cdot) \\ \infty & \text{otherwise;} \end{cases}$$

(ii) for every  $x \in U$ ,  $\widehat{L}(x, \cdot)$  is the lsc envelope of  $L(x, \cdot)$ .



## 5. PROOF OF THEOREM 2.1 AND 2.2

5.1. **Proof of Theorem 2.1.** Let  $O \in \mathcal{O}(\Omega)$  and  $u \in \text{dom}F(\cdot, O)$ . If  $\mathcal{Z}f$  is ru-usc and  $W^{1,p}$ -quasiconvex, then by Lemma 4.2 we have

$$\overline{F}(u, O) \geq \inf \left\{ \varliminf_{n \rightarrow \infty} \int_O \mathcal{Z}f(x, \nabla u_n(x)) dx : W^{1,p}(\Omega; \mathbb{R}^m) \ni u_n \rightharpoonup u \right\}.$$

Using Proposition 2.1 (ii) we obtain

$$(5.1) \quad \overline{F}(u, O) \geq \int_O \widehat{\mathcal{Z}f}(x, \nabla u(x)) dx.$$

If  $f$  is ru-usc then (5.1) holds by Proposition 2.1 (iii).

To prove the reverse inequality, note that  $tu \in \text{dom}F(\cdot, O)$  for all  $t \in ]0, 1[$  since (2.5) and (2.3). Using Lemma 4.2 and (2.5) we have for every  $t \in ]0, 1[$

$$\mathcal{Z}f(x, t\nabla u(x)) \leq f(x, t\nabla u(x)) \leq C(1 + f(x, 0) + f(x, \nabla u(x))) \text{ a.e. in } O$$

Then considering both Proposition 2.1 (i) and Lemma 4.8 and applying the Lebesgue dominated theorem we obtain

$$\begin{aligned} \overline{F}(u, O) &\leq \varliminf_{t \rightarrow 1^-} \overline{F}(tu, O) \leq \overline{\lim}_{t \rightarrow 1^-} \overline{F}(tu, O) \leq \overline{\lim}_{t \rightarrow 1^-} \int_O \mathcal{Z}f(x, t\nabla u) dx \\ &\leq \int_O \widehat{\mathcal{Z}f}(x, \nabla u) dx, \end{aligned}$$

where we used the fact that  $\overline{F}(\cdot, O)$  is swlsc on  $W^{1,p}(\Omega; \mathbb{R}^m)$ , which completes the proof.  $\blacksquare$

5.2. **Proof of Theorem 2.2.** Fix  $O \in \mathcal{O}(\Omega)$ . By (2.11) and (2.5)  $\text{dom}F(\cdot, O)$  is a convex subset of  $W^{1,p}(\Omega; \mathbb{R}^m)$  with 0 belongs to the interior of  $\text{dom}F(\cdot, O)$  with respect the norm topology of  $W^{1,p}(\Omega; \mathbb{R}^m)$ . Hence, by a well known property of convex set in normed space, we have  $t\overline{\text{dom}F(\cdot, O)}^s \subset \text{dom}F(\cdot, O)$  for all  $t \in [0, 1[$ , where  $\overline{\text{dom}F(\cdot, O)}^s$  is the closure of  $\text{dom}F(\cdot, O)$  in  $W^{1,p}(\Omega; \mathbb{R}^m)$ . Since  $\text{dom}F(\cdot, O)$  is a convex set we have  $\overline{\text{dom}F(\cdot, O)}^s = \overline{\text{dom}F(\cdot, O)}^w$  where  $\overline{\text{dom}F(\cdot, O)}^w$  is the closure of  $\text{dom}F(\cdot, O)$  with respect to the weak topology of  $W^{1,p}(\Omega; \mathbb{R}^m)$ . We deduce that  $\text{dom}F(\cdot, O)$  is weakly star shaped with respect to 0, i.e.,

$$(5.2) \quad t\overline{\text{dom}F(\cdot, O)}^w \subset \text{dom}F(\cdot, O) \text{ for all } t \in [0, 1[.$$

We claim that  $\overline{F}(\cdot, O)$  is ru-usc in  $\text{dom}F(\cdot, O)$ . Indeed, let  $u \in \text{dom}F(\cdot, O)$  and  $t \in ]0, 1[$ . First, by Proposition 2.1 (i)

$$\begin{aligned} \overline{F}(tu, O) &\leq \int_O \widehat{\mathcal{Z}f}(x, t\nabla u) dx \leq \int_O \Delta_{\widehat{\mathcal{Z}f}}^a(t) \left( a(x) + \widehat{\mathcal{Z}f}(x, \nabla u) \right) + \widehat{\mathcal{Z}f}(x, \nabla u) dx \\ &= \Delta_{\widehat{\mathcal{Z}f}}^a(t) (|a|_{L^1(O)} + \overline{F}(u, O)) + \overline{F}(u, O). \end{aligned}$$

It follows that  $\Delta_{\overline{F}(\cdot, O), \text{dom}F(\cdot, O)}^{|a|_{L^1(O)}}(t) \leq \Delta_{\widehat{\mathcal{Z}f}}^a(t)$  for all  $t \in ]0, 1[$ . By Lemma 4.8 (if  $f$  is ru-usc then combine Lemma 4.7 and Lemma 4.8 and  $(\text{dom}\widehat{\mathcal{Z}f} \subset \overline{\text{dom}\mathcal{Z}f})$   $\widehat{\mathcal{Z}f}$  is ru-usc, it follows that  $\overline{F}(\cdot, O)$  is ru-usc in  $\text{dom}F(\cdot, O)$ . Applying Corollary 4.2 with  $I(u) = \int_O \widehat{\mathcal{Z}f}(x, \nabla u) dx$ ,  $D = \text{dom}F(\cdot, O)$  and by taking account of (5.2), we obtain

$$(5.3) \quad \overline{F}(u, O) := \begin{cases} \int_O \widehat{\mathcal{Z}f}(x, \nabla u) dx & \text{if } u \in \text{dom}F(\cdot, O) \\ \varliminf_{t \rightarrow 1} \int_O \widehat{\mathcal{Z}f}(x, t\nabla u) dx & \text{if } u \in \text{dom}\overline{F}(\cdot, O) \setminus \text{dom}F(\cdot, O) \\ \infty & \text{otherwise.} \end{cases}$$

Let  $u \in \text{dom}\bar{F}(\cdot, O) \setminus \text{dom}F(\cdot, O)$ . If  $I(u) = \infty$  then  $\bar{F}(u, O) = \infty$ , indeed, since

$$\liminf_{t \rightarrow 1} \liminf_{s \rightarrow 1} \mathcal{Z}f(x, st\xi) \geq \widehat{\mathcal{Z}f}(x, \xi),$$

we have by (5.3) and Fatou lemma

$$\bar{F}(u, O) = \lim_{t \rightarrow 1} \int_O \widehat{\mathcal{Z}f}(x, t\nabla u) dx \geq \int_O \liminf_{t \rightarrow 1} \widehat{\mathcal{Z}f}(x, t\nabla u) dx \geq \int_O \widehat{\mathcal{Z}f}(x, \nabla u) dx = I(u).$$

Assume now that  $I(u) < \infty$ . On one hand,  $\widehat{\mathcal{Z}f}(\cdot, \nabla u(\cdot)) \in L^1(O)$ , and on the other hand  $\widehat{\mathcal{Z}f}$  is ru-usc, hence

$$\widehat{\mathcal{Z}f}(x, t\nabla u(x)) \leq \widehat{\mathcal{Z}f}(x, \nabla u(x)) + \Delta_{\widehat{\mathcal{Z}f}}^a(t) \left( a(x) + \widehat{\mathcal{Z}f}(x, \nabla u(x)) \right)$$

for all  $t \in ]0, 1[$  and  $x \in O$ . Applying the Lebesgue dominated theorem we finally obtain

$$\lim_{t \rightarrow 1} \int_O \widehat{\mathcal{Z}f}(x, t\nabla u) dx = \int_O \widehat{\mathcal{Z}f}(x, \nabla u) dx. \quad \blacksquare$$

## 6. PROOF OF PROPOSITION 2.1 (i)

**6.1. Local Dirichlet problems associated to a functional.** For any functional  $H : W^{1,p}(\Omega; \mathbb{R}^m) \times \mathcal{O}(\Omega) \rightarrow [0, \infty]$  we set

$$m_H(u; O) := \inf \left\{ H(v; O) : v \in u + W_0^{1,p}(O; \mathbb{R}^m) \right\}.$$

Note that we can write also  $m_H(u; O) = \inf \left\{ H(u + \varphi; O) : \varphi \in W_0^{1,p}(O; \mathbb{R}^m) \right\}$ . For each  $\varepsilon > 0$  and each  $O \in \mathcal{O}(\Omega)$ , denote by  $\mathcal{V}_\varepsilon(O)$  the class of all countable family  $\{\bar{Q}_i := \bar{Q}_{\rho_i}(x_i)\}_{i \in I}$  of disjoint (pairwise disjoint) closed balls of  $O$  with  $x_i \in O$  and  $\rho_i = \text{diam}(Q_i) \in ]0, \varepsilon[$  such that  $|O \setminus \cup_{i \in I} Q_i| = 0$ . Consider  $m_H^\varepsilon(u; \cdot) : \mathcal{O}(\Omega) \rightarrow [0, \infty]$  given by

$$m_H^\varepsilon(u; O) := \inf \left\{ \sum_{i \in I} m_H(u; Q_i) : \{\bar{Q}_i\}_{i \in I} \in \mathcal{V}_\varepsilon(O) \right\},$$

and define  $m_H^*(u; \cdot) : \mathcal{O}(\Omega) \rightarrow [0, \infty]$  by

$$m_H^*(u; O) := \sup_{\varepsilon > 0} m_H^\varepsilon(u; O) = \lim_{\varepsilon \rightarrow 0} m_H^\varepsilon(u; O).$$

The set function  $m_H^*$  is of the Carathéodory construction type (see for instance [Fed69, 2.10]), it was introduced by [BFM98] and [BB00b].

**Lemma 6.1.** *Let  $O \in \mathcal{O}(\Omega)$ . Assume that  $H(u; \cdot)$  is countably subadditive for all  $u \in W^{1,p}(O; \mathbb{R}^m)$ . Then for every  $u \in W^{1,p}(O; \mathbb{R}^m)$  we have*

$$(6.1) \quad m_H(u; O) \leq m_H^*(u; O).$$

*Proof.* Fix  $\varepsilon > 0$ . Choose  $\{\bar{Q}_i\}_{i \geq 1} \in \mathcal{V}_\varepsilon(O)$  such that

$$(6.2) \quad \sum_{i \geq 1} m_H(u; Q_i) \leq \frac{\varepsilon}{2} + m_H^*(u; O).$$

For each  $i \geq 1$  there exists  $\varphi_\varepsilon^i \in W_0^{1,p}(Q_i; \mathbb{R}^m)$  such that

$$(6.3) \quad H(u + \varphi_\varepsilon^i; Q_i) \leq \frac{\varepsilon}{2^{i+1}} + m_H(u; Q_i).$$

Set  $\varphi_\varepsilon := \sum_{i \geq 1} \varphi_\varepsilon^i \mathbb{I}_{Q_i} \in W_0^{1,p}(O; \mathbb{R}^m)$ . Using the countable subadditivity of  $H(u; \cdot)$ , (6.3), and (6.2) we have

$$\begin{aligned} m_H(u; O) &\leq H(u + \varphi_\varepsilon; O) \leq \sum_{i \geq 1} H(u + \varphi_\varepsilon^i; Q_i) \leq \frac{\varepsilon}{2} + \sum_{i \geq 1} m_H(u; Q_i) \\ &\leq \varepsilon + m_H^*(u; O), \end{aligned}$$

we obtain (6.1) by letting  $\varepsilon \rightarrow 0$ .  $\blacksquare$

By [BB00b, Prop. 2.1., p. 81], we have the following result (which is needed for the proof of Lemma 6.3).

**Lemma 6.2.** *Let  $u \in W^{1,p}(\Omega; \mathbb{R}^m)$ . If there exists a finite Radon measure  $\mu_u$  on  $\Omega$  such that for every cube  $Q \in \mathcal{O}(\Omega)$*

$$m_H(u; Q) \leq \mu_u(Q),$$

*then  $m_H^*(u; \cdot)$  can be extended to a Radon measure  $\lambda_u$  on  $\Omega$  satisfying  $0 \leq \lambda_u \leq \mu_u$ .*

The proof of the upper bound will be divided into four steps.

6.2.  $\bar{F}(\mathbf{u}; \mathbf{O}) \leq \mathbf{m}_F^*(\mathbf{u}; \mathbf{O})$  for all  $(\mathbf{u}, \mathbf{O}) \in \mathbf{W}^{1,p}(\Omega; \mathbb{R}^m) \times \mathcal{O}(\Omega)$ . Fix  $u \in W^{1,p}(\Omega; \mathbb{R}^m)$ . Without loss of generality we assume that  $m_F^*(u; O) < \infty$ . Let  $\varepsilon \in ]0, 1[$ . There exists  $\{\bar{Q}_i\}_{i \in I} \in \mathcal{V}_\varepsilon(O)$  such that

$$(6.4) \quad \sum_{i \in I} m_F(u; Q_i) \leq m_F^\varepsilon(u; O) + \frac{\varepsilon}{2} \leq m_F^*(u; O) + \frac{\varepsilon}{2}.$$

Given any  $i \in I$ , by definition of  $m_F(u; Q_i)$ , there exists  $v_i \in u + W_0^{1,p}(Q_i; \mathbb{R}^m)$  such that

$$(6.5) \quad F(v_i; Q_i) \leq m_F^\varepsilon(u; Q_i) + \frac{\varepsilon}{2} \frac{|Q_i|}{|O|}.$$

Define  $u_\varepsilon \in u + W_0^{1,p}(O; \mathbb{R}^m)$  by  $u_\varepsilon := \sum_{i \in I} v_i \mathbb{I}_{Q_i} + u \mathbb{I}_{\Omega \setminus \cup_{i \in I} Q_i}$ . From (6.4) and (6.5) we have that

$$(6.6) \quad F(u_\varepsilon; O) \leq m_F^\varepsilon(u; O) + \varepsilon.$$

In the case  $p \in ]d, \infty[$ , from the  $p$ -coercivity of  $f$ , (6.4) and (6.5), we deduce

$$(6.7) \quad \sup_{\varepsilon > 0} \int_O |\nabla u_\varepsilon|^p dx \leq \frac{1}{c} (m_F^*(u; O) + 1).$$

By Poincaré inequality there exists  $K > 0$  depending on  $p$  and  $d$  only such that for each  $v_i \in u + W_0^{1,p}(Q_i; \mathbb{R}^m)$

$$\int_{Q_i} |v_i - u|^p dx \leq K \varepsilon^p \int_{Q_i} |\nabla v_i - \nabla u|^p dx,$$

since  $\text{diam}(Q_i) < \varepsilon$ . By summing on  $i \in I$  and using (6.7) and we obtain

$$\begin{aligned} \int_O |u_\varepsilon - u|^p dx &\leq 2^{p-1} K \varepsilon^p \left( \int_O |\nabla u_\varepsilon|^p dx + \int_O |\nabla u|^p dx \right) \\ &\leq 2^{p-1} K \varepsilon^p \left( \frac{1}{c} (m_F^*(u; O) + 1) + \int_O |\nabla u|^p dx \right) \end{aligned}$$

which shows that  $u_\varepsilon \rightarrow u$  in  $L^p(O; \mathbb{R}^m)$  as  $\varepsilon \rightarrow 0$ . In the case where  $p = \infty$ , from (6.6) we have

$$(6.8) \quad \|\nabla u_\varepsilon\|_{L^\infty(O; \mathbb{R}^m)} \leq R_0$$

and reasoning similarly as in the case  $p$  finite we obtain  $u_\varepsilon \rightarrow u$  in  $L^\infty(O; \mathbb{R}^m)$  as  $\varepsilon \rightarrow 0$ .

Therefore by (6.7) ((6.8) if  $p = \infty$ ), there is a subsequence (not relabeled) such that  $u_\varepsilon \rightharpoonup u$  ( $u_\varepsilon \xrightarrow{*} u$  if  $p = \infty$ ) as  $\varepsilon \rightarrow 0$ , and then by (6.6) we have

$$\overline{F}(u; O) \leq \varliminf_{\varepsilon \rightarrow 0} \overline{F}(u_\varepsilon; O) \leq \mathbf{m}_F^*(u; O).$$

■

*Remark 6.1.* We note that the previous proof establishes

$$\begin{aligned} \overline{F}(u; O) &\leq \inf \left\{ \varliminf_{\varepsilon \rightarrow 0} F(u_\varepsilon; O) : u + W_0^{1,p}(O; \mathbb{R}^m) \ni u_\varepsilon \rightharpoonup u \text{ in } W^{1,p}(\Omega; \mathbb{R}^m) \right\} \\ &\leq \mathbf{m}_F^*(u; O). \end{aligned}$$

**6.3.  $\mathbf{m}_F^*(u; \cdot)$  is locally equivalent to  $\mathbf{m}_F(u; \cdot)$ .** We are concerned with the proof of the local equivalence of  $\mathbf{m}_F^*(u; \cdot)$  and  $\mathbf{m}_F(u; \cdot)$ , this result was established by [BFM98, Lemma 3.5] in the context of relaxation of variational functionals in  $BV$ , and in a general framework in [BB00b, Theorem 2.3]. However, the proof that we propose is inspired by [ABF03, Proof of Theorem 3.11, p. 380]. Note that by Lemma 6.1  $\varliminf_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F^*(u; \mathbf{Q}_\varepsilon(x_0))}{\mathbf{m}_F(u; \mathbf{Q}_\varepsilon(x_0))} \geq 1$ .

**Lemma 6.3.** *If  $F(u; O) < \infty$ . Then we have*

$$\lim_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F^*(u; \mathbf{Q}_\varepsilon(x_0))}{\varepsilon^d} = \lim_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F(u; \mathbf{Q}_\varepsilon(x_0))}{\varepsilon^d} \quad \text{a.e. in } O.$$

*Proof.* Let  $u \in W^{1,p}(\Omega; \mathbb{R}^m)$  be such that  $F(u; O) < \infty$ . Then for each  $U \in \mathcal{O}(O)$

$$\mathbf{m}_F(u; U) \leq \int_U f(x, \nabla u(x)) dx < \infty,$$

so by using Lemma 6.2 with  $\mu_u := f(\cdot, \nabla u(\cdot)) dx|_O$ ,  $\mathbf{m}_F^*(u; \cdot)$  is the trace of a Radon measure  $\lambda_u$  on  $O$  satisfying  $0 \leq \lambda_u \leq \mu_u$ . Since  $\mu_u$  is absolutely continuous with respect to  $dx|_O$  the Lebesgue measure on  $O$ , the limit  $\lim_{\varepsilon \rightarrow 0} \frac{\lambda_u(\mathbf{Q}_\varepsilon(x_0))}{\varepsilon^d}$  exists for a.a.  $x_0 \in O$  as the Radon-Nikodym derivative of  $\lambda_u$  with respect to  $dx|_O$ . Moreover, by Lemma 8.6 we have

$$\lim_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F^*(u; \mathbf{Q}_\varepsilon(x_0))}{\varepsilon^d} \geq \varliminf_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F(u; \mathbf{Q}_\varepsilon(x_0))}{\varepsilon^d} \quad \text{a.e. in } O.$$

It remains to prove that

$$(6.9) \quad \lim_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F^*(u; \mathbf{Q}_\varepsilon(x_0))}{\varepsilon^d} \leq \varliminf_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F(u; \mathbf{Q}_\varepsilon(x_0))}{\varepsilon^d} \quad \text{a.e. in } O.$$

Fix any  $\theta > 0$ . Consider the following sets

$$\begin{aligned} \mathcal{G}_\theta &:= \left\{ \mathbf{Q}_\varepsilon(x) : x \in O, \varepsilon > 0 \text{ and } \mathbf{m}_F^*(u; \mathbf{Q}_\varepsilon(x)) > \mathbf{m}_F(u; \mathbf{Q}_\varepsilon(x)) + \theta |\mathbf{Q}_\varepsilon(x)| \right\}, \\ \mathcal{N}_\theta &:= \left\{ x \in O : \forall \delta > 0 \exists \varepsilon \in ]0, \delta[ \mathbf{Q}_\varepsilon(x) \in \mathcal{G}_\theta \right\}. \end{aligned}$$

It is sufficient to prove that  $\mathcal{N}_\theta$  is a negligible set for the Lebesgue measure on  $O$ , indeed, in this case given  $x_0 \in O \setminus \mathcal{N}_\theta$  there exists  $\delta_0 > 0$  such that  $\mathbf{m}_F^*(u; \mathbf{Q}_\varepsilon(x_0)) \leq \mathbf{m}_F(u; \mathbf{Q}_\varepsilon(x_0)) + \theta |\mathbf{Q}_\varepsilon(x_0)|$  for all  $\varepsilon \in ]0, \delta_0[$ . Hence

$$\lim_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F^*(u; \mathbf{Q}_\varepsilon(x_0))}{|\mathbf{Q}_\varepsilon(x_0)|} \leq \varliminf_{\varepsilon \rightarrow 0} \frac{\mathbf{m}_F(u; \mathbf{Q}_\varepsilon(x_0))}{|\mathbf{Q}_\varepsilon(x_0)|} + \theta,$$

then by letting  $\theta \rightarrow 0$  we obtain (6.9).

Fix  $\delta > 0$ . Consider the set

$$\mathcal{F}_\delta := \left\{ \overline{\mathbf{Q}}_\varepsilon(x) : x \in \mathcal{N}_\theta, \varepsilon \in ]0, \delta[ \text{ and } \mathbf{Q}_\varepsilon(x) \in \mathcal{G}_\theta \right\}.$$

Using the definition of  $\mathcal{N}_\theta$  we can see that  $\inf_{Q \in \mathcal{F}_\delta} \text{diam}(Q) = 0$ . By the Vitali covering theorem there exists a disjointed countable subfamily  $\{\overline{Q}_i\}_{i \geq 1}$  of  $\mathcal{F}_\delta$  such that

$$(6.10) \quad |\mathcal{N}_\theta \setminus \bigcup_{i \geq 1} Q_i| = 0.$$

We have  $\mathcal{N}_\theta \subset \bigcup_{i \geq 1} Q_i \cup \mathcal{N}_\theta \setminus \bigcup_{i \geq 1} Q_i$ , then to prove that  $\mathcal{N}_\theta$  is a negligible set, it is equivalent to prove that  $|V_j| = 0$  for all  $j \geq 1$ , where

$$V_j := \bigcup_{i=1}^j Q_i.$$

Fix  $j \geq 1$ . Let  $\{Q'_i\}_{i \geq 1} \in \mathcal{V}_\delta(O \setminus \bigcup_{i=1}^j \overline{Q}_i)$  satisfying

$$(6.11) \quad \sum_{i \geq 1} m_F(u; Q'_i) \leq m_F^*(u; O \setminus \bigcup_{i=1}^j \overline{Q}_i) + \delta.$$

Recalling that  $m_F^*(u; \cdot)$  is the trace on  $\mathcal{O}(O)$  of a nonnegative finite Radon measure, we see that

$$\begin{aligned} m_F^*(u; O) &\geq m_F^*(u; O \setminus \bigcup_{i=1}^j \overline{Q}_i) + m_F^*(u; V_j) \\ &= m_F^*(u; O \setminus \bigcup_{i=1}^j \overline{Q}_i) + \sum_{1 \leq i \leq j} m_F^*(u; Q_i). \end{aligned}$$

Since each  $Q_i \in \mathcal{G}_\theta$ , we have by using (6.11)

$$m_F^*(u; O) \geq \sum_{i \geq 1} m_F(u; Q'_i) - \delta + \sum_{i=1}^j m_F(u; Q_i) + \theta |V_j|.$$

It is easy to see that the countable family  $\{Q'_i : i \geq 1\} \cup \{Q_i : 1 \leq i \leq j\}$  belongs to  $\mathcal{V}_\delta(O)$ , thus

$$m_F^*(u; O) \geq m_F^\delta(u; O) + \theta |V_j| - \delta.$$

Letting  $\delta \rightarrow 0$  we have  $m_F^\delta(u; O) \nearrow m_F^*(u; O)$ , and so  $|V_j| = 0$  since  $\theta > 0$ .  $\blacksquare$

**6.4. Cut-off technique to substitute  $u(\cdot)$  with  $u(x_0) + \nabla u(x_0)(\cdot - x_0)$  in  $\mathbf{m}_F(u; \cdot)$ .** Here using cut-off functions we show that for almost all  $x_0 \in \Omega$  we can (locally) replace  $u$  in  $m_F(u; \cdot)$  with the affine tangent map of  $u$  at  $x_0$  denoted by  $u_{x_0}(\cdot) := u(x_0) + \nabla u(x_0)(\cdot - x_0)$ . In the following, we consider  $u \in W^{1,p}(\Omega; \mathbb{R}^m)$  satisfying  $tu \in \text{dom}F$  for all  $t \in ]0, 1[$ .

We claim that for every  $t \in ]0, 1[$

$$(6.12) \quad \lim_{\varepsilon \rightarrow 0} \frac{m_F(tu; Q_\varepsilon(x_0))}{\varepsilon^d} \leq \mathcal{Z}f(x_0, t\nabla u(x_0)) \quad \text{a.e. } x_0 \in \Omega.$$

Fix  $t \in ]0, 1[$  and consider  $\lambda, \alpha \in ]0, 1[$  such that  $\lambda = \frac{t}{\alpha}$ . Fix  $r, s \in ]0, 1[$  such that  $s < r$ . Fix  $x_0 \in \Omega$  such that

$$(6.13) \quad \lim_{\varepsilon \rightarrow 0} \frac{F(\alpha u; Q_\varepsilon(x_0))}{\varepsilon^d} = f(x_0, \alpha \nabla u(x_0)) < \infty;$$

$$(6.14) \quad \lim_{\varepsilon \rightarrow 0} \frac{F(tu; Q_\varepsilon(x_0))}{\varepsilon^d} = f(x_0, t\nabla u(x_0)) < \infty;$$

$$(6.15) \quad \mathcal{M}_0(x_0) = \lim_{\varepsilon \rightarrow 0} \int_{Q_\varepsilon(x_0)} \mathcal{M}_0(x) dx < \infty.$$

$$(6.16) \quad \mathcal{Z}f(x_0, t\nabla u(x_0)) = \lim_{\varepsilon \rightarrow 0} \inf_{\varphi \in W_0^{1,p}(Q_\varepsilon(x_0); \mathbb{R}^m)} \int_{Q_\varepsilon(x_0)} f(y, t\nabla u(x_0) + \nabla \varphi) dy.$$

To shorten notation we denote by  $Q_\varepsilon$  the cube  $Q_\varepsilon(x_0)$ .

Let  $\{\varepsilon_n\}_n \subset \mathbb{R}_+^*$  be a sequence such that  $\varepsilon_n \rightarrow 0$  as  $n \rightarrow \infty$  and

$$\lim_{\varepsilon \rightarrow 0} \frac{\mathfrak{m}_F(tu; \mathbb{Q}_\varepsilon)}{\varepsilon^d} = \lim_{n \rightarrow \infty} \frac{\mathfrak{m}_F(tu; \mathbb{Q}_{\varepsilon_n})}{\varepsilon_n^d}.$$

Fix  $n \geq 1$ . Choose  $v_n \in u_{x_0} + W_0^{1,p}(\mathbb{Q}_{s\varepsilon_n}; \mathbb{R}^m)$  such that

$$F(tv_n; \mathbb{Q}_{s\varepsilon_n}) \leq \mathfrak{m}_F(tu_{x_0}; \mathbb{Q}_{s\varepsilon_n}) + (\varepsilon_n)^{d+1},$$

Consider a cut-off function  $\phi \in W_0^{1,\infty}(\mathbb{Q}_{\varepsilon_n}; [0, 1])$  such that  $\|\nabla\phi\|_{L^\infty(\mathbb{Q}_{\varepsilon_n})} \leq \frac{4}{(r-s)\varepsilon_n}$  and

$$\phi(x) = \begin{cases} 1 & \text{on } \mathbb{Q}_{s\varepsilon_n} \\ 0 & \text{on } \mathbb{Q}_{\varepsilon_n} \setminus \mathbb{Q}_{r\varepsilon_n} \end{cases}, \quad [0 < \phi < 1] \subset\subset \mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}.$$

Define  $w_n := \phi v_n + (1 - \phi)u \in u + W_0^{1,p}(\mathbb{Q}_{\varepsilon_n}; \mathbb{R}^m)$ , we have

(6.17)

$$\begin{aligned} \mathfrak{m}_F(tu; \mathbb{Q}_{\varepsilon_n}) &\leq F(tv_n; \mathbb{Q}_{s\varepsilon_n}) + F(tw_n; \mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}) + F(tu; \mathbb{Q}_{\varepsilon_n} \setminus \mathbb{Q}_{r\varepsilon_n}) \\ &\leq \mathfrak{m}_F(tu_{x_0}; \mathbb{Q}_{s\varepsilon_n}) + \varepsilon_n^{d+1} + F(tw_n; \mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}) + F(tu; \mathbb{Q}_{\varepsilon_n} \setminus \mathbb{Q}_{r\varepsilon_n}). \end{aligned}$$

The rest of the proof consists to give estimates from above, as  $n \rightarrow \infty$ , of the two last terms of (6.17) divided by  $\varepsilon_n^d$ .

By (6.14) we have

(6.18)

$$\lim_{n \rightarrow \infty} \frac{F(tu; \mathbb{Q}_{\varepsilon_n} \setminus \mathbb{Q}_{r\varepsilon_n})}{\varepsilon_n^d} = \lim_{n \rightarrow \infty} \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{\varepsilon_n} \setminus \mathbb{Q}_{r\varepsilon_n}} f(x, t\nabla u) dx = (1 - r^d) f(x_0, t\nabla u(x_0)).$$

By (2.5) we have

(6.19)

$$\begin{aligned} &\frac{F(tw_n; \mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n})}{\varepsilon_n^d} \\ &= \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} f(x, \lambda(\psi\alpha\nabla u(x_0) + (1 - \psi)\alpha\nabla u) + (1 - \lambda)\Phi_{n,t}) dx \\ &\leq C_1 \left( \frac{|\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}|}{\varepsilon_n^d} + \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} f(x, \alpha\nabla u(x_0)) dx \right. \\ &\quad \left. + \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} f(x, \alpha\nabla u) dx + \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} f(x, \Phi_{n,t}) dx \right), \end{aligned}$$

where  $\Phi_{n,t} := \frac{t}{1-\lambda} \nabla\phi \otimes (u_{x_0} - u)$  and  $C_1 = C^2 + C$ . By (6.13), it holds

(6.20)

$$\lim_{n \rightarrow \infty} \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} f(x, \alpha\nabla u) dx = (r^d - s^d) f(x_0, \alpha\nabla u(x_0)),$$

Using (2.4) and (6.13) we deduce

(6.21)

$$\lim_{n \rightarrow \infty} \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} f(x, \alpha\nabla u(x_0)) dx = (r^d - s^d) f(x_0, \alpha\nabla u(x_0)).$$

Choose  $N_0 \geq 1$ , such that  $\frac{1}{\varepsilon_n} \|u_{x_0} - u\|_{L^\infty(\mathbb{Q}_{\varepsilon_n})} \leq \frac{(1-\lambda)(r-s)\rho_0}{4t}$  for all  $n \geq N_0$ . It follows that  $\|\Phi_{n,t}\|_{L^\infty(\mathbb{Q}_{\varepsilon_n}; \mathbb{M}^{m,d})} \leq \rho_0$  for all  $n \geq N_0$ , and by Lemma 4.1

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} f(x, \Phi_{n,t}) dx \leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} \mathcal{M}_0(x) dx.$$

Moreover, by (6.15) we have

(6.22)

$$\overline{\lim}_{n \rightarrow \infty} \frac{1}{\varepsilon_n^d} \int_{\mathbb{Q}_{r\varepsilon_n} \setminus \mathbb{Q}_{s\varepsilon_n}} \mathcal{M}_0(x) dx = (r^d - s^d) \mathcal{M}_0(x_0).$$

Passing to the limit  $n \rightarrow \infty$  by taking account of (6.19), and the estimates (6.20), (6.21), (6.22), we have

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{\mathfrak{m}_F(tu; \mathbb{Q}_{\varepsilon_n})}{\varepsilon_n^d} \\ & \leq s^d \lim_{n \rightarrow \infty} \frac{\mathfrak{m}_F(tu_{x_0}; \mathbb{Q}_{s\varepsilon_n})}{s^d \varepsilon_n^d} + (1 - r^d) f(x_0, t\nabla u(x_0)) \\ & \quad + 2C_1(r^d - s^d)(1 + f(x_0, \alpha\nabla u(x_0)) + f(x_0, \alpha\nabla u(x_0))) + \mathcal{M}_0(x_0). \end{aligned}$$

Letting  $r \rightarrow 1$  and  $s \rightarrow 1$ , we find

$$(6.23) \quad \lim_{\varepsilon \rightarrow 0} \frac{\mathfrak{m}_F(tu; \mathbb{Q}_\varepsilon)}{\varepsilon^d} = \lim_{n \rightarrow \infty} \frac{\mathfrak{m}_F(tu; \mathbb{Q}_{\varepsilon_n})}{\varepsilon_n^d} \leq \lim_{s \rightarrow 1} \lim_{n \rightarrow \infty} \frac{\mathfrak{m}_F(tu_{x_0}; \mathbb{Q}_{s\varepsilon_n})}{(s\varepsilon_n)^d}$$

$$(6.24) \quad \begin{aligned} & \leq \overline{\lim}_{\varepsilon \rightarrow 0} \frac{\mathfrak{m}_F(tu_{x_0}; \mathbb{Q}_\varepsilon)}{\varepsilon^d} \\ & = \mathcal{Z}f(x_0, t\nabla u(x_0)), \end{aligned}$$

where we have used (6.16).

**6.5. End of the proof of Proposition 2.1 (i).** Using in turn the results of Subsect. 6.2, 6.3 and 6.4 we obtain for every  $u \in W^{1,p}(\Omega; \mathbb{R}^m)$ , every  $O \in \mathcal{O}(\Omega)$  and every  $t \in ]0, 1[$

$$\begin{aligned} \overline{F}(tu; O) & \leq \mathfrak{m}_F^*(tu; O) = \int_O \lim_{\varepsilon \rightarrow 0} \frac{\mathfrak{m}_F^*(tu; \mathbb{Q}_\varepsilon(x))}{\varepsilon^d} dx = \int_O \lim_{\varepsilon \rightarrow 0} \frac{\mathfrak{m}_F(tu; \mathbb{Q}_\varepsilon(x))}{\varepsilon^d} dx \\ & \leq \int_O \mathcal{Z}f(x, t\nabla u(x)) dx. \end{aligned}$$

■

## 7. PROOF OF PROPOSITION 2.1 (ii) AND (iii)

The proof will be divided into two steps. The first one is a localization technique also known as blow-up method introduced by [FM92] which consists to reduce the proof of the (global) lower bound to a local lower bound by using measure arguments. The second step consists to proof the local lower bound by using cut-off functions.

In this section we denote by  $L$  the integrands  $\mathcal{Z}f$  or  $f$ .

**7.1. Localization technique.** Let  $O \in \mathcal{O}(\Omega)$ . Let  $u, \{u_n\}_n \subset W^{1,p}(\Omega; \mathbb{R}^m)$  be such that  $u_n \rightharpoonup u$  in  $W^{1,p}(\Omega; \mathbb{R}^m)$  and

$$\begin{aligned} \infty > \mathcal{L}(u; O) & := \inf \left\{ \liminf_{n \rightarrow \infty} \int_O L(x, \nabla u_n(x)) dx : W^{1,p}(\Omega; \mathbb{R}^m) \ni u_n \rightharpoonup u \right\} \\ & = \lim_{n \rightarrow \infty} \int_O L(x, \nabla u_n(x)) dx. \end{aligned}$$

Up to a subsequence, since  $p > d$ , we may assume that

$$(7.1) \quad u_n \rightarrow u \text{ in } L^\infty(\Omega; \mathbb{R}^m).$$

Passing to a subsequence if necessary, we may find a nonnegative Radon measure  $\mu$  such that

$$L(\cdot, \nabla u_n(\cdot)) dx|_O \xrightarrow{*} \mu \text{ as } n \rightarrow \infty \text{ weakly } * \text{ in the sense of measures.}$$

It is enough to prove that for all  $t \in ]0, 1[$

$$(7.2) \quad \frac{d\mu}{dx}(x_0) + \Delta_L^a(t) \left( a(x_0) + \frac{d\mu}{dx}(x_0) \right) \geq \mathcal{Z}L(x_0, t\nabla u(x_0)) \quad x_0 \text{ a.e. in } O.$$

Indeed, by Alexandrov theorem, we will have

$$\mathcal{L}(u; O) = \lim_{n \rightarrow \infty} L(u_n; O) \geq \varliminf_{n \rightarrow \infty} \int_O L(x, \nabla u_n) dx = \mu(O) \geq \int_O \frac{d\mu}{dx}(x) dx,$$

so by integrating over  $O$  in (7.2), we find

$$\mathcal{L}(u; O) + \Delta_L^a(t) (|a|_{L^1(O)} + \mathcal{L}(u; O)) \geq \int_O \mathcal{Z}L(x, t\nabla u(x)) dx.$$

Since  $L$  is ru-usc, we obtain the result by passing to the limit  $t \rightarrow 1$  and by using Fatou lemma.

As  $\int_\Omega f(x, \nabla u(x)) dx < \infty$ , we fix  $x_0 \in O$  such that (2.4) holds and

$$(7.3) \quad f(x_0, \nabla u(x_0)) < \infty,$$

$$(7.4) \quad L(x_0, \nabla u(x_0)) \leq f(x_0, \nabla u(x_0)) < \infty,$$

$$(7.5) \quad \frac{d\mu}{dx}(x_0) = \lim_{\varepsilon \rightarrow 0} \frac{\mu(Q_\varepsilon(x_0))}{\varepsilon^d} < \infty,$$

$$(7.6) \quad \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \|u - u(x_0) - \nabla u(x_0) \cdot (\cdot - x_0)\|_{L^\infty(Q_\varepsilon(x_0); \mathbb{R}^m)} = 0,$$

$$(7.7) \quad a(x_0) = \lim_{\varepsilon \rightarrow 0} \int_{Q_\varepsilon(x_0)} a(x) dx < \infty,$$

$$(7.8) \quad \mathcal{M}_0(x_0) = \lim_{\varepsilon \rightarrow 0} \int_{Q_\varepsilon(x_0)} \mathcal{M}_0(x) dx < \infty,$$

where  $Q_\varepsilon(x_0) := x_0 + \varepsilon Y$ . Note that (7.8) is a consequence of Lemma 4.1.

Choose  $\varepsilon_k \rightarrow 0$  such that  $\mu(\partial Q_{\varepsilon_k}(x_0)) = 0$ . Then

$$(7.9) \quad \begin{aligned} \lim_{k \rightarrow \infty} \frac{\mu(Q_{\varepsilon_k}(x_0))}{\varepsilon_k^d} &= \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{Q_{\varepsilon_k}(x_0)} L(x, \nabla u_n) dx \\ &= \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \int_Y L(x_0 + \varepsilon_k y, \nabla v_{n,k}) dy, \end{aligned}$$

where  $v_{n,k}(y) := \frac{u_n(x_0 + \varepsilon_k y) - u(x_0)}{\varepsilon_k}$ . By (7.1) we have

$$(7.10) \quad \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \|v_{n,k} - l_{\nabla u(x_0)}\|_{L^\infty(Y; \mathbb{R}^m)} = 0 \quad \text{where} \quad l_{\nabla u(x_0)}(y) := \nabla u(x_0)y.$$

Fix  $s, r \in ]0, 1[$  such that  $s < r$ . Then (7.5) implies (see subsect. 7.2.1 for the proof)

$$(7.11) \quad \lim_{k \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \frac{\mu_n(Q_{r\varepsilon_k}(x_0) \setminus \overline{Q_{s\varepsilon_k}(x_0)})}{\varepsilon_k^d} = (r^d - s^d) \frac{d\mu}{dx}(x_0).$$

By a simultaneous diagonalization of (7.9), (7.10), (7.11), we may extract a subsequence  $v_n := v_{n, k_n}$  satisfying

$$(7.12) \quad v_n \rightarrow l_{\nabla u(x_0)} \text{ in } L^\infty(Y; \mathbb{R}^m), \quad v_n \rightharpoonup l_{\nabla u(x_0)} \text{ in } W^{1,p}(Y; \mathbb{R}^m),$$

$$(7.13) \quad \frac{d\mu}{dx}(x_0) = \lim_{n \rightarrow \infty} \int_Y L(x_0 + \varepsilon_n y, \nabla v_n) dy,$$

$$(7.14) \quad (r^d - s^d) \frac{d\mu}{dx}(x_0) = \lim_{n \rightarrow \infty} \frac{\mu_n(Q_{r\varepsilon_n}(x_0) \setminus \overline{Q_{s\varepsilon_n}(x_0)})}{\varepsilon_n^d},$$

where  $\varepsilon_{k_n} := \varepsilon_n$ .



**7.2. Cut-off technique to substitute  $v_n$  with  $w_n \in l_{\nabla u(x_0)} + W^{1,p}(Y; \mathbb{R}^m)$ .** For simplicity of notation we set  $\theta_{x_0,n}(y) := x_0 + \varepsilon_n y$  for all  $y \in Y$ . In this section we use cut-off functions to show that there exists  $\{w_n\}_n \subset l_{\nabla u(x_0)} + W_0^{1,p}(Y; \mathbb{R}^m)$  such that for every  $t \in ]0, 1[$

$$(7.15) \quad \overline{\lim}_{n \rightarrow \infty} \int_Y L(\theta_{x_0,n}, t \nabla w_n) dy \leq \frac{d\mu}{dx}(x_0) + \Delta_L^a(t) \left( a(x_0) + \frac{d\mu}{dx}(x_0) \right).$$

If (7.15) holds then

$$\begin{aligned} \mathcal{Z}L(x_0, t \nabla u(x_0)) &\leq \liminf_{\varepsilon \rightarrow 0} \left\{ \int_Y L(x_0 + \varepsilon y, t \nabla w) dy : w \in l_{\nabla u(x_0)} + W_0^{1,p}(Y; \mathbb{R}^m) \right\} \\ &\leq \overline{\lim}_{n \rightarrow \infty} \int_Y L(\theta_{x_0,n}, t \nabla w_n) dy \\ &= \frac{d\mu}{dx}(x_0) + \Delta_L^a(t) \left( a(x_0) + \frac{d\mu}{dx}(x_0) \right), \end{aligned}$$

and the claim 7.2 follows.

So, let us prove (7.15). Fix any  $t \in ]0, 1[$ . Let  $\phi \in W_0^{1,\infty}(Y; [0, 1])$  be a cut-off function between  $s\bar{Y}$  and  $\bar{Y} \setminus rY$  such that  $\|\nabla \phi\|_{L^\infty(Y)} \leq \frac{4}{r-s}$ . Setting

$$w_n := \phi v_n + (1 - \phi) l_{\nabla u(x_0)}.$$

We have  $w_n \in l_{\nabla u(x_0)} + W_0^{1,p}(Y; \mathbb{R}^m)$  and

$$\nabla w_n := \begin{cases} \nabla v_n & \text{on } sY \\ \phi \nabla v_n + (1 - \phi) \nabla u(x_0) + \Phi_{n,s,r} & \text{on } U_{s,r} \\ \nabla u(x_0) & \text{on } Y \setminus r\bar{Y}, \end{cases}$$

where  $\Phi_{n,s,r} := \nabla \phi \otimes (v_n - l_{\nabla u(x_0)})$  and  $U_{s,r} := rY \setminus s\bar{Y}$ .

For every  $n \geq 1$ , it holds

$$(7.16) \quad \begin{aligned} &\int_Y L(\theta_{x_0,n}, t \nabla w_n) dy \\ &= \int_{sY} L(\theta_{x_0,n}, t \nabla v_n) dy + \int_{U_{s,r}} L(\theta_{x_0,n}, t \nabla w_n) dy + \int_{Y \setminus r\bar{Y}} L(\theta_{x_0,n}, t \nabla u(x_0)) dy \\ &\leq \int_Y L(\theta_{x_0,n}, t \nabla v_n) dy + \int_{U_{s,r}} L(\theta_{x_0,n}, t \nabla w_n) dy + \int_{Y \setminus r\bar{Y}} L(\theta_{x_0,n}, t \nabla u(x_0)) dy. \end{aligned}$$

The rest of the proof consists to give estimates from above, as  $n \rightarrow \infty$ , of the three last terms of (7.16).

**Bound for  $\overline{\lim}_{n \rightarrow \infty} \int_Y L(\theta_{x_0,n}, t \nabla v_n) dy$ .** Since  $L$  is ru-usc, using (7.13) and (7.7), we have for every  $n \geq 1$

$$(7.17) \quad \begin{aligned} &\overline{\lim}_{n \rightarrow \infty} \int_Y L(\theta_{x_0,n}, t \nabla v_n) dy \\ &\leq \overline{\lim}_{n \rightarrow \infty} \left( \Delta_L^a(t) \int_Y a(\theta_{x_0,n}) + L(\theta_{x_0,n}, \nabla v_n) dy + \int_Y L(\theta_{x_0,n}, \nabla v_n) dy \right) \\ &\leq \Delta_L^a(t) \left( a(x_0) + \frac{d\mu}{dx}(x_0) \right) + \frac{d\mu}{dx}(x_0). \end{aligned}$$

**Bound for  $\overline{\lim}_{n \rightarrow \infty} \int_{Y \setminus r\overline{Y}} L(\theta_{x_0, n}, t \nabla u(x_0)) dy$ .** Similarly to the previous estimate we have

$$\begin{aligned} & \overline{\lim}_{n \rightarrow \infty} \int_{Y \setminus r\overline{Y}} L(\theta_{x_0, n}, t \nabla u(x_0)) dy \\ & \leq \overline{\lim}_{n \rightarrow \infty} \left( \Delta_L^a(t) \int_{Y \setminus r\overline{Y}} a(\theta_{x_0, n}) + L(\theta_{x_0, n}, \nabla u(x_0)) dy + \int_{Y \setminus r\overline{Y}} L(\theta_{x_0, n}, \nabla u(x_0)) dy \right) \\ & \leq \Delta_L^a(t) ((1 - r^d)a(x_0) + A_r(x_0)) + A_r(x_0), \end{aligned}$$

where  $A_r(x_0) := \overline{\lim}_{n \rightarrow \infty} \int_{Y \setminus r\overline{Y}} L(\theta_{x_0, n}, \nabla u(x_0)) dy$ . Now, taking account of (7.3) and (7.4), by (2.4) we have an upper bound for  $A_r(x_0)$

$$(7.18) \quad A_r(x_0) \leq (1 - r^d)f(x_0, \nabla u(x_0)).$$

We deduce

$$(7.19) \quad \begin{aligned} & \overline{\lim}_{n \rightarrow \infty} \int_{Y \setminus r\overline{Y}} L(\theta_{x_0, n}, t \nabla u(x_0)) dy \\ & \leq (1 - r^d) (\Delta_L^a(t)(a(x_0) + L(x_0, \nabla u(x_0))) + f(x_0, \nabla u(x_0))). \end{aligned}$$

**Bound for  $\overline{\lim}_{n \rightarrow \infty} \int_{U_{s,r}} L(\theta_{x_0, n}, t \nabla w_n) dy$ .** It remains to estimate the middle term of (7.16). Since  $f$  satisfies (2.5), by Lemma 4.3 we have that  $L = \mathcal{Z}f$  also satisfies (2.5). Therefore for every  $n \geq 1$

$$\begin{aligned} & \int_{U_{s,r}} L(\theta_{x_0, n}, t \nabla w_n) dy \\ & \leq C_1 \left( (r^d - s^d) + \int_{U_{s,r}} L(\theta_{x_0, n}, \nabla v_n) dy + \int_{U_{s,r}} L(\theta_{x_0, n}, \nabla u(x_0)) dy \right. \\ & \quad \left. + \int_{U_{s,r}} L \left( \theta_{x_0, n}, \frac{t}{1-t} \Phi_{n,s,r} \right) dy \right) \end{aligned}$$

Since (7.12), there exists  $n_{s,r} \geq 1$  such that for every  $n \geq n_{s,r}$

$$\left\| \frac{t}{1-t} \Phi_{n,s,r} \right\|_{L^\infty(Y; \mathbb{M}^{m \times d})} \leq \rho_0,$$

where  $\rho_0 > 0$  is given by Lemma 4.1. Taking account of (7.8), we have

$$(7.20) \quad \begin{aligned} \overline{\lim}_{n \rightarrow \infty} \int_{U_{s,r}} L \left( \theta_{x_0, n}, \frac{t}{1-t} \Phi_{n,s,r} \right) dy & \leq \overline{\lim}_{n \rightarrow \infty} \int_{U_{s,r}} \sup_{\zeta \in \overline{Q}_{\rho_0}(0)} L(\theta_{x_0, n}, \zeta) dy \\ & \leq (r^d - s^d) \mathcal{M}_0(x_0). \end{aligned}$$

Reasoning similarly to the estimate (7.18), we find

$$(7.21) \quad \overline{\lim}_{n \rightarrow \infty} \int_{U_{s,r}} L(\theta_{x_0, n}, \nabla u(x_0)) dy \leq (r^d - s^d) f(x_0, \nabla u(x_0)).$$

Since (7.14), we have

$$(7.22) \quad \overline{\lim}_{n \rightarrow \infty} \int_{U_{s,r}} L(\theta_{x_0, n}, \nabla v_n) dy = (r^d - s^d) \frac{d\mu}{dx}(x_0).$$

Collecting (7.20), (7.21) and (7.22), we obtain

$$(7.23) \quad \begin{aligned} & \overline{\lim}_{n \rightarrow \infty} \int_{U_{s,r}} L(\theta_{x_0, n}, t \nabla w_n) dy \\ & \leq C_1 (s^d - r^d) \left( 1 + \frac{d\mu}{dx}(x_0) + f(x_0, \nabla u(x_0)) + \mathcal{M}_0(x_0) \right). \end{aligned}$$

**End of the proof of (7.15).** Collecting (7.17), (7.19) and (7.23), we have

$$\begin{aligned}
& \overline{\lim}_{n \rightarrow \infty} \int_Y L(\theta_{x_0, n}, t \nabla w_n) dy \\
& \leq \Delta_L^a(t) \left( a(x_0) + \frac{d\mu}{dx}(x_0) \right) + \frac{d\mu}{dx}(x_0) \\
& + (1 - r^d) (\Delta_L^a(t) (a(x_0) + L(x_0, \nabla u(x_0))) + f(x_0, \nabla u(x_0))) \\
& + C_1 (r^d - s^d) \left( 1 + \frac{d\mu}{dx}(x_0) + f(x_0, \nabla u(x_0)) + \mathcal{M}_0(x_0) \right).
\end{aligned}$$

Letting  $r \rightarrow 1$  and  $s \rightarrow 1$  we obtain (7.15). ■

7.2.1. *Proof of (7.11).* By (7.5) we have

$$(7.24) \quad \frac{d\mu}{dx}(x_0) = \lim_{\varepsilon \rightarrow 0} \frac{\mu(Q_{s\varepsilon}(x_0))}{(s\varepsilon)^d} = \lim_{\varepsilon \rightarrow 0} \frac{\mu(Q_{r\varepsilon}(x_0))}{(r\varepsilon)^d} = \lim_{\varepsilon \rightarrow 0} \frac{\mu(\overline{Q}_{s\varepsilon}(x_0))}{(s\varepsilon)^d} = \lim_{\varepsilon \rightarrow 0} \frac{\mu(\overline{Q}_{r\varepsilon}(x_0))}{(r\varepsilon)^d}.$$

Using (7.24) and Alexandrov theorem on one hand we have

$$(7.25) \quad \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \frac{\mu_n(Q_{r\varepsilon_k}(x_0) \setminus \overline{Q}_{s\varepsilon_k}(x_0))}{\varepsilon_k^d} \geq \lim_{k \rightarrow \infty} \frac{\mu(Q_{r\varepsilon_k}(x_0) \setminus \overline{Q}_{s\varepsilon_k}(x_0))}{\varepsilon_k^d} = (r^d - s^d) \frac{d\mu}{dx}(x_0).$$

Similarly, on the other hand we have

$$(7.26) \quad \begin{aligned} & \overline{\lim}_{k \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \frac{\mu_n(Q_{r\varepsilon_k}(x_0) \setminus \overline{Q}_{s\varepsilon_k}(x_0))}{\varepsilon_k^d} \\ & \leq \overline{\lim}_{k \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \left( r^d \frac{\mu_n(Q_{r\varepsilon_k}(x_0))}{(r\varepsilon_k)^d} - s^d \frac{\mu_n(\overline{Q}_{s\varepsilon_k}(x_0))}{(s\varepsilon_k)^d} \right) \\ & \leq \overline{\lim}_{k \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \left( r^d \frac{\mu_n(\overline{Q}_{r\varepsilon_k}(x_0))}{(r\varepsilon_k)^d} - s^d \frac{\mu_n(Q_{s\varepsilon_k}(x_0))}{(s\varepsilon_k)^d} \right) \\ & \leq \overline{\lim}_{k \rightarrow \infty} \left( r^d \frac{\mu(\overline{Q}_{r\varepsilon_k}(x_0))}{(r\varepsilon_k)^d} - s^d \frac{\mu(Q_{s\varepsilon_k}(x_0))}{(s\varepsilon_k)^d} \right) = (r^d - s^d) \frac{d\mu}{dx}(x_0) \end{aligned}$$

Combining (7.25) and (7.26), we obtain (7.11). ■

## 8. PROOF OF PROPOSITION 2.2

We denote by  $\text{Cub}$  the family of all open cubes of  $\mathbb{R}^d$ . We denote by  $\text{Cub}_\delta$  the family of all open cube  $Q$  of  $\mathbb{R}^d$  such that  $\text{diam}(Q) < \delta$ , where  $\delta > 0$ . For each  $E \subset \mathbb{R}^d$ , we associate the set  $\mathcal{F}_\delta(E)$  of all countable families  $\{Q_i\}_{i \in I} \subset \text{Cub}_\delta$  satisfying  $|E \setminus \cup_{i \in I} Q_i| = 0$ ,  $Q_i \cap E \neq \emptyset$  for all  $i \in I$ , and  $\overline{Q}_i \cap \overline{Q}_j = \emptyset$  for all  $i \neq j$ . If  $E \neq \emptyset$  then  $\mathcal{F}_\delta(E) \neq \emptyset$ , indeed, by the Vitali covering theorem, it is always possible starting from a family of closed cubes of  $\mathbb{R}^d$  with center in  $E$  to find a countable subfamily of open cubes in  $\mathcal{F}_\delta(E)$  because the Lebesgue measure of the boundary of a cube is null, i.e.,  $|\overline{Q}| = |Q|$  for all  $Q \in \text{Cub}$ .

Let  $m$  be a nonnegative set function defined for all cube of  $\mathbb{R}^d$  such that  $m(\emptyset) = 0$ . Let  $m^\sharp : \mathcal{P}(E) \rightarrow [0, \infty]$  be defined by

$$m^\sharp(E) := \begin{cases} \sup_{\delta>0} m^\delta(E) & \text{if } E \neq \emptyset \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{with } m^\delta(E) := \inf \left\{ \sum_{i \in I} m(Q_i) : \{Q_i\}_{i \in I} \in \mathcal{F}_\delta(E) \right\}.$$

We denote by  $\omega \in [0, \infty]$  the number

$$\omega := \overline{\lim}_{\delta \rightarrow 0} \sup_{\substack{Q \subset \Omega \\ \text{diam}(Q) < \delta}} \frac{m(Q)}{|Q|}$$

where  $Q$  denotes any arbitrary open cube of  $\mathbb{R}^d$ .

The following result is an abstract version of Proposition 2.2.

**Proposition 8.1.** *If  $\omega < \infty$  and*

$$(8.1) \quad \overline{\lim}_{\delta \rightarrow 0} \frac{m(Q_\delta(x))}{\delta^d} \leq \overline{\lim}_{\delta \rightarrow 0} \frac{m^\sharp(Q_\delta(x))}{\delta^d} \quad \text{a.e. in } \Omega$$

then

$$\overline{\lim}_{\delta \rightarrow 0} \frac{m(Q_\delta(x))}{\delta^d} = \lim_{\delta \rightarrow 0} \frac{m(Q_\delta(x))}{\delta^d} \quad \text{a.e. in } \Omega,$$

where  $Q_\delta(x) = x + \delta Y$  for any  $x \in \Omega$  and  $\delta > 0$ .

The set function  $m^\sharp$  is of Carathéodory type construction [Fed69, Sect. 2.10, p. 169], but we do not know whether it is an outer measure, however we have the following result.

**Lemma 8.1.** *The set function  $m^\sharp$  satisfies:*

- (i) *if  $E_1, E_2 \subset \mathbb{R}^d$  are two sets such that  $\text{dist}(E_1, E_2) > 0$  then  $m^\sharp(E_1 \cup E_2) = m^\sharp(E_1) + m^\sharp(E_2)$ ;*
- (ii) *if  $E, V \subset \mathbb{R}^d$  are such that  $V$  is a nonempty open set and  $E \subset V$  then  $m^\sharp(E) \leq m^\sharp(V)$ ;*
- (iii) *if  $\omega < \infty$  then  $m^\sharp(E) \leq \omega|E|$  for all closed set  $E \subset \Omega$ .*

*Proof.* (i) We show that for every  $E_1, E_2 \subset \mathbb{R}^d$  satisfying  $\text{dist}(E_1, E_2) > \delta_0$  for some  $\delta_0 > 0$  we have

$$(8.2) \quad m^\sharp(E_1 \cup E_2) \geq m^\sharp(E_1) + m^\sharp(E_2).$$

Fix  $\delta \in ]0, \delta_0[$ . Choose  $\{Q_i\}_{i \in I} \in \text{Cub}_\delta$  satisfying  $|(E_1 \cup E_2) \setminus \cup_{i \in I} Q_i| = 0$ ,  $Q_i \cap (E_1 \cup E_2) \neq \emptyset$  for all  $i \in I$ , and

$$(8.3) \quad m^\sharp(E_1 \cup E_2) + \delta \geq \sum_{i \in I} m(Q_i).$$

Let  $I_j = \{i \in I : Q_i \cap E_j \neq \emptyset\}$  for  $j \in \{1, 2\}$ . Since  $\text{dist}(E_1, E_2) > 2\delta$ , if  $i \in I_1$  (resp.  $i \in I_2$ ) then  $Q_i \cap E_2 = \emptyset$  (resp.  $Q_i \cap E_1 = \emptyset$ ). Thus

$$0 = |(E_1 \cup E_2) \setminus \cup_{i \in I} Q_i| = |E_1 \setminus \cup_{i \in I_1} Q_i| + |E_2 \setminus \cup_{i \in I_2} Q_i|,$$

hence  $|E_j \setminus \cup_{i \in I_j} Q_i| = 0$  and  $Q_i \cap E_j \neq \emptyset$  for all  $i \in I_j$ . From (8.3) we have

$$m^\sharp(E_1 \cup E_2) + \delta \geq \sum_{i \in I_1} m(Q_i) + \sum_{i \in I_2} m(Q_i) \geq m^\sharp(E_1) + m^\sharp(E_2),$$

and (8.2) holds by letting  $\delta \rightarrow 0$ .

Now, we show that

$$(8.4) \quad m^\sharp(E_1 \cup E_2) \leq m^\sharp(E_1) + m^\sharp(E_2).$$

For each  $j \in \{1, 2\}$ , choose  $\{Q_i^j\}_{i \in I_j} \in \text{Cub}_\delta$  satisfying  $|E_j \setminus \bigcup_{i \in I_j} Q_i^j| = 0$ ,  $Q_i^j \cap E_j \neq \emptyset$  for all  $i \in I_j$ , and

$$(8.5) \quad m^\sharp(E_j) + \delta \geq \sum_{i \in I_j} m(Q_i^j).$$

Since  $\text{dist}(E_1, E_2) > \delta_0$  the countable family of cubes  $\{Q_i^j : i \in I_j \text{ and } j \in \{1, 2\}\}$  is pairwise disjoint, moreover we have

$$|(E_1 \cup E_2) \setminus \bigcup_{i \in I_1} Q_i \cup \bigcup_{i \in I_2} Q_i| \leq |E_1 \setminus \bigcup_{i \in I_1} Q_i| + |E_2 \setminus \bigcup_{i \in I_2} Q_i| = 0.$$

Summing over  $j \in \{1, 2\}$  in (8.5) we obtain

$$m^\sharp(E_1) + m^\sharp(E_2) + 2\delta \geq \sum_{j \in \{1, 2\}} \sum_{i \in I_j} m(Q_i^j) \geq m^\sharp(E_1 \cup E_2),$$

and (8.4) holds by letting  $\delta \rightarrow 0$ .

(ii) Let  $E, V$  be two sets of  $\mathbb{R}^d$  such that  $V$  is a nonempty open set and  $E \subset V$ . For each  $\delta > 0$  choose  $\{Q_i\}_{i \in I} \in \text{Cub}_\delta$  satisfying  $|V \setminus \bigcup_{i \in I} Q_i| = 0$ ,  $Q_i \cap V \neq \emptyset$  for all  $i \in I$ , and

$$(8.6) \quad m^\sharp(V) + \delta \geq \sum_{i \in I} m(Q_i).$$

Consider the open set  $V_\delta := \bigcup_{i \in I} Q_i$ , then  $|V \setminus \overline{V_\delta}| = 0$ , but  $V \setminus \overline{V_\delta}$  is open so  $V \setminus \overline{V_\delta} = \emptyset$ . Which means that  $V \subset \overline{V_\delta}$ , therefore  $V \subset V_\delta$ . We deduce that  $I_E \neq \emptyset$  where  $I_E := \{i \in I : Q_i \cap E \neq \emptyset\}$ . We have  $|E \setminus \bigcup_{i \in I_E} Q_i| = |E \setminus \bigcup_{i \in I} Q_i| \leq |V \setminus \bigcup_{i \in I} Q_i| = 0$ , thus  $\{Q_i\}_{i \in I_E} \in \mathcal{F}_\delta(E)$  and so from (8.6)

$$m^\sharp(V) + \delta \geq m^\sharp(E),$$

and (ii) holds by letting  $\delta \rightarrow 0$ .

(iii) Fix  $\delta > 0$  and  $E \subset \Omega$ . Set  $E_\delta = \{x \in \mathbb{R}^d : \text{dist}(x, E) < \delta\}$ , then for any countable family  $\{Q_i\}_{i \in I} \in \mathcal{F}_\delta(E)$  we have  $\bigcup_{i \in I} Q_i \subset E_\delta$  since  $Q_i \cap E \neq \emptyset$  for all  $i \in I$  and  $\text{diam}(Q) < \delta$ . Therefore

$$\begin{aligned} m^\delta(E) &\leq \sum_{i \in I} m(Q_i) \leq \sum_{i \in I} \frac{m(Q_i)}{|Q_i|} |Q_i| \leq \sup_{\text{diam}(Q) < \delta} \frac{m(Q)}{|Q|} \left| \bigcup_{i \in I} Q_i \right| \\ &\leq \sup_{\text{diam}(Q) < \delta} \frac{m(Q)}{|Q|} |E_\delta|. \end{aligned}$$

Passing to limit  $\delta \rightarrow 0$  we obtain  $m^\sharp(E) \leq \omega |E|$ . ■

Let  $m_*^+, m_*^- : \Omega \rightarrow [0, \infty]$  be the functions defined by

$$m_*^+(x) := \overline{\lim}_{\delta \rightarrow 0} \sup_{x \in Q \in \text{Cub}_\delta} \frac{m^\sharp(Q)}{|Q|} \quad \text{and} \quad m_*^-(x) := \underline{\lim}_{\delta \rightarrow 0} \inf_{x \in Q \in \text{Cub}_\delta} \frac{m(Q)}{|Q|},$$

**Lemma 8.2.** *Let  $a, b \in \mathbb{R}^+$ . Then*

- (i) *there exists a Borel set  $B_a^+ \subset \{x \in \Omega : m_*^+(x) \geq a\}$  such that  $|\{x \in \Omega : m_*^+(x) \leq a\} \setminus B_a^+| = 0$ ;*
- (ii) *there exists a Borel set  $B_b^- \subset \{x \in \Omega : m_*^-(x) \leq b\}$  such that  $|\{x \in \Omega : m_*^-(x) \leq a\} \setminus B_b^-| = 0$ .*

*Proof.* Let us prove (ii). Let  $b \in \mathbb{R}^+$ . Set  $M_b = \{x \in \Omega : m_*^-(x) \leq b\}$ . For each  $k \in \mathbb{N}^*$ , consider the set

$$\mathcal{G}_k := \{\overline{Q}_\varepsilon(x) : x \in M_b \text{ and } \varepsilon \in ]0, \frac{1}{k}[ \text{ and } m(Q_\varepsilon(x)) \leq (b + \frac{1}{k})|Q_\varepsilon(x)|\}$$

For each  $k \geq 1$  the family  $\mathcal{G}_k$  is a fine cover of  $M_b$ , and so by the Vitali covering theorem, there exists a disjointed countable family  $\{\overline{Q}_i^k\}_{i \in I_k} \subset \mathcal{G}_k$  such that  $|M \setminus \cup_{i \in I_k} Q_i^k| = 0$ . Consider the Borel set  $B_b^- := \cap_{k \geq 1} \cup_{i \in I_k} Q_i^k$ , then  $|M_b \setminus B_b^-| \leq \sum_{k \geq 1} |M_b \setminus \cup_{i \in I_k} Q_i^k| = 0$ . If we show that  $B_b^- \subset M_b$  then the proof of (ii) will be complete. Let  $y \in B_b^-$ . Then for every  $k \geq 1$  there exists  $i_k \in I_k$  such that  $y \in Q_{i_k}^k \in \text{Cub}_{\frac{1}{k}}$  and  $m(Q_{i_k}^k) \leq (b + \frac{1}{k})|Q_{i_k}^k|$ . It follows that

$$\inf_{y \in Q \in \text{Cub}_{\frac{1}{k}}} \frac{m(Q)}{|Q|} \leq \frac{m(Q_{i_k}^k)}{|Q_{i_k}^k|} \leq b + \frac{1}{k},$$

letting  $k \rightarrow \infty$ , we obtain that  $m_*^-(y) \leq b$  which means that  $y \in M_b$ .

For the proof of (i), it is enough to remark that for  $a > 0$

$$\{x \in \Omega : m_*^+(x) \geq a\} = \left\{ x \in \Omega : \lim_{\delta \rightarrow 0} \inf_{x \in Q \in \text{Cub}_\delta} \frac{|Q|}{m^\sharp(Q)} \leq \frac{1}{a} \right\},$$

and to apply the same reasoning as in the proof of (ii) with the necessary changes.  $\blacksquare$

*Remark 8.1.* By Lemma 8.2, the functions  $m_*^-$  and  $m_*^+$  are measurable.

*Remark 8.2.* The same conclusions can be drawn if we replace large inequalities with strict inequalities in the Lemma 8.2, indeed, it suffices to see for instance that

$$\{x \in \Omega : m_*^+(x) > a\} = \bigcup_{n \geq 1} \{x \in \Omega : m_*^+(x) \geq a + \frac{1}{n}\}.$$

We denote by  $\overline{m}^\sharp$  the set function

$$\overline{m}^\sharp(E) = \inf \{m^\sharp(O) : E \subset O, O \text{ open}\} \text{ for all } E \subset \mathbb{R}^d.$$

**Lemma 8.3.** *If  $\omega < \infty$  then  $\overline{m}^\sharp(K) = m^\sharp(K)$  for all compact  $K \subset \Omega$ .*

*Proof.* Fix a compact set  $K \subset \Omega$ . Note that by Lemma 8.1 (ii) we have  $\overline{m}^\sharp(K) \geq m^\sharp(K)$ . So it remains to prove the reverse inequality  $\overline{m}^\sharp(K) \leq m^\sharp(K)$ .

By Lemma 8.1 (iii) we have  $m^\sharp(K) \leq \omega|K| \leq \omega|\Omega| < \infty$ . Let  $O \subset \Omega$  be an open set such that  $O \supset K$ . For each  $n \in \mathbb{N}^*$  such that  $n \geq n_0$  where  $n_0 := \text{Ent}((\text{diam}(O) - \text{diam}(K))^{-1}) + 1$  ( $\text{Ent}(r)$  denotes the integer part of the real number  $r$ ) there exists  $\{Q_j^n\}_{j \geq 1} \subset \mathcal{F}_{\frac{1}{n}}(K)$  such that

$$\infty > m^\sharp(K) + \frac{1}{n} \geq \sum_{j \geq 1} m(Q_j^n).$$

Note that  $\cup_{j \geq 1} Q_j^n \subset \{x \in \mathbb{R}^d : \text{dist}(x, K) < \frac{1}{n}\} \subset O$  because  $n \geq n_0$  and  $Q_j^n \cap K \neq \emptyset$  for all  $j \geq 1$ .

Fix  $n \geq n_0$ . Then there exists an increasing sequence  $\{j_s\}_{s \geq 1}$  such that  $\sup_{s \geq 1} j_s = \infty$  and  $\alpha_s := \sum_{j \geq j_s} m(Q_j^n) \leq \frac{1}{s}$  for all  $s \geq 1$ . Fix  $s \in \mathbb{N}^*$ . For the open set  $O \setminus \cup_{1 \leq j \leq j_s} \overline{Q}_j^n$  we use the Vitali covering theorem to find a disjointed countable family of closed cubes  $\{\overline{Q}_i^n\}_{i \in I}$  such that  $\text{diam}(Q_i^n) < \frac{1}{n}$ ,

$$\left| \left( O \setminus \bigcup_{1 \leq j \leq j_s} \overline{Q}_j^n \right) \setminus \bigcup_{i \in I} \overline{Q}_i^n \right| = 0, \text{ and } \bigcup_{i \in I} \overline{Q}_i^n \subset O \setminus \bigcup_{1 \leq j \leq j_s} \overline{Q}_j^n.$$

It is direct to see that the countable family

$$\{\overline{Q}_k^n\}_{k \in D} := \{\overline{Q}_i^n : i \in I\} \cup \{\overline{Q}_j^n : 1 \leq j \leq j_s\} \in \mathcal{F}_{\frac{1}{n}}(O).$$

If  $\omega_{\frac{1}{n}} := \sup_{Q \subset \Omega, \text{diam}(Q) < \frac{1}{n}} \frac{m(Q)}{|Q|}$  then

$$\begin{aligned} m^{\frac{1}{n}}(O) - m^{\sharp}(K) &\leq \sum_{k \in D} m(Q_k^n) - \sum_{j \geq 1} m(Q_j^n) + \frac{1}{n} \\ &= \sum_{i \in I} m(Q_i^n) - \alpha_s + \frac{1}{n} \\ &\leq \omega_{\frac{1}{n}} |O \setminus \bigcup_{1 \leq j \leq j_s} \overline{Q}_j^n| - \alpha_s + \frac{1}{n}. \end{aligned}$$

Passing to the limit  $s \rightarrow \infty$  we obtain  $m^{\frac{1}{n}}(O) - m^{\sharp}(K) \leq \omega_{\frac{1}{n}} |O \setminus \bigcup_{j \geq 1} \overline{Q}_j^n| + \frac{1}{n}$ . If  $E := \bigcap_{n \geq n_0} \bigcup_{j \geq 1} Q_j^n$  then  $|K \setminus E| = 0$ , indeed, we have

$$|K \setminus E| \leq \sum_{n \geq n_0} |K \setminus \bigcup_{j \geq 1} \overline{Q}_j^n| = 0.$$

Letting  $n \rightarrow \infty$  it follows that  $m^{\sharp}(O) - m^{\sharp}(K) \leq \omega |O \setminus E|$ . Therefore

$$\begin{aligned} m^{\sharp}(O) &\leq m^{\sharp}(K) + \omega (|(O \setminus K) \setminus E| + |K \setminus E|) \\ &\leq m^{\sharp}(K) + \omega |O \setminus K|. \end{aligned}$$

Since the open set  $O$  containing  $K$  is arbitrary, by the outer regularity of the Lebesgue measure we obtain  $\overline{m^{\sharp}}(K) \leq m^{\sharp}(K)$ , and the proof is complete.  $\blacksquare$

**Lemma 8.4.** *Let  $a, b > 0$ . Let  $E \subset \Omega$  be an arbitrary set.*

- (i) *If  $E \subset \{x \in \Omega : m_*^+(x) > a\}$  then  $\overline{m^{\sharp}}(E) \geq a|E|$ ;*
- (ii) *If  $E \subset \{x \in \Omega : m_*^-(x) < b\}$  then  $m^{\sharp}(E) \leq b|E|$ .*

*Proof.* We start by the proof of (i). Fix  $a > 0$  and let  $E \subset \{x \in \Omega : m_*^+(x) > a\}$ . Let  $O$  be an open set of  $\Omega$  such that  $O \supset E$ . We can rewrite

$$\{x \in \Omega : m_*^+(x) > a\} = \left\{ x \in \Omega : \lim_{\delta \rightarrow 0} \inf_{x \in Q \in \text{Cub}_{\delta}} \frac{|Q|}{m^{\sharp}(Q)} < \frac{1}{a} \right\}.$$

Fix  $\delta > 0$  and consider the family of closed cubes

$$\mathcal{G}_{\delta} := \{\overline{Q}_{\varepsilon}(x) : x \in E, \text{Cub}_{\delta} \ni Q_{\varepsilon}(x) \subset O \text{ and } |Q_{\varepsilon}(x)| \leq \frac{1}{a} m^{\sharp}(Q_{\varepsilon}(x))\}.$$

The family  $\mathcal{G}_{\delta}$  is a fine covering of  $E$ . By the Vitali covering theorem, there exists a disjointed countable family  $\{\overline{Q}_i\}_{i \in I} \subset \mathcal{G}_{\delta}$  such that  $|E \setminus \bigcup_{i \in I} Q_i| = 0$ . For each  $\varepsilon > 0$  there exists a finite set  $I_{\varepsilon} \subset I$  such that  $|E \setminus \bigcup_{i \in I_{\varepsilon}} Q_i| < \varepsilon$ . Then by using Lemma 8.1 (i)

$$|E| - \varepsilon = |E \cap \bigcup_{i \in I_{\varepsilon}} Q_i| \leq \sum_{i \in I_{\varepsilon}} |Q_i| \leq \frac{1}{a} \sum_{i \in I_{\varepsilon}} m^{\sharp}(Q_i) = m^{\sharp}(\bigcup_{i \in I_{\varepsilon}} Q_i) \leq m^{\sharp}(O).$$

The proof of (i) is complete since the open set  $O$  which contains  $E$  is arbitrary.

It remains to prove (ii). For each  $\delta > 0$  consider the set

$$\mathcal{G}_{\delta} := \{\overline{Q}_{\varepsilon}(x) : Q_{\varepsilon}(x) \in \text{Cub}_{\delta}, x \in E \text{ and } m(Q_{\varepsilon}(x)) \leq b|Q_{\varepsilon}(x)|\}.$$

It is a fine cover of  $E$ , i.e.,  $\inf\{\text{diam}(Q) : Q \in \mathcal{G}_{\delta}\} = 0$ . Then there exists a disjointed countable subfamily  $\{Q_i\}_{i \in I} \subset \mathcal{G}_{\delta}$  such that  $|E \setminus \bigcup_{i \in I} Q_i| = 0$  and  $\sum_{i \in I} |Q_i| \leq |E| + \delta$  (see [Mat95, Theorem 2.2, p. 26]), so  $\{Q_i\}_{i \in I} \in \mathcal{F}_{\delta}(E)$ . It follows that

$$m^{\sharp}(E) \leq \sum_{i \in I} m(Q_i) \leq \sum_{i \in I} b|Q_i| \leq b|E| + b\delta,$$

and the proof of (ii) is complete by letting  $\delta \rightarrow 0$ .  $\blacksquare$

**Lemma 8.5.** *If  $\omega < \infty$  then  $m_*^+(x) \leq m_*^-(x)$  a.e. in  $\Omega$ .*

*Proof.* Fix  $a, b \in \mathbb{Q}$  such that  $a > b > 0$ . Consider the following set

$$S_{a,b} := \{x \in \Omega : m_*^-(x) < b < a < m_*^+(x)\}.$$

By Remark 8.2 and Lemma 8.2 there exists a Borel set  $B_{a,b}$  such that  $B_{a,b} \subset S_{a,b}$  and  $|S_{a,b} \setminus B_{a,b}| = 0$ . Fix  $\varepsilon > 0$ . Since the Lebesgue measure is inner regular, choose a compact set  $K_\varepsilon \subset B_{a,b}$  such that  $|B_{a,b} \setminus K_\varepsilon| < \varepsilon$ . From Lemma 8.3 we have  $\overline{m}^\#(K_\varepsilon) = m^\#(K_\varepsilon)$  since  $\omega < \infty$ . Using Lemma 8.4 we obtain  $a|K_\varepsilon| \leq m^\#(K_\varepsilon) \leq b|K_\varepsilon|$ . Therefore  $|K_\varepsilon| = 0$  since  $b < a$ , hence  $|S_{a,b}| = |B_{a,b} \setminus K_\varepsilon| + |K_\varepsilon| + |S_{a,b} \setminus B_{a,b}| < \varepsilon$ , and so by letting  $\varepsilon \rightarrow 0$  we find  $|S_{a,b}| = 0$ . Now, the set where  $m_*^+$  is greater than  $m_*^-$  is a countable union of negligible sets, i.e.,

$$\{x \in \Omega : m_*^-(x) < m_*^+(x)\} = \bigcup_{0 < b < a, (a,b) \in \mathbb{Q}^2} S_{a,b},$$

and the proof is complete.  $\blacksquare$

**Proof of Proposition 8.1.** Using (8.1) and the definitions of  $m_*^+$  and  $m_*^-$  we have for every  $x \in \Omega$

$$m_*^-(x) \leq \liminf_{\delta \rightarrow 0} \frac{m(Q_\delta(x))}{\delta^d} \leq \overline{\lim}_{\delta \rightarrow 0} \frac{m(Q_\delta(x))}{\delta^d} \leq m_*^+(x).$$

By Lemma 8.5 we obtain

$$m_*^-(x) = m_*^+(x) = \lim_{\delta \rightarrow 0} \frac{m(Q_\delta(x))}{\delta^d} \text{ a.e. in } \Omega.$$

which completes the proof.  $\blacksquare$

If  $L : \Omega \times \mathbb{M}^{m \times d} \rightarrow [0, \infty]$  is a Borel measurable integrand then for each  $\xi \in \mathbb{M}^{m \times d}$  we denote by  $m_\xi : \text{Cub} \rightarrow [0, \infty]$  the set function defined by

$$m_\xi(Q) = \inf \left\{ \int_Q L(x, \xi + \nabla \varphi(x)) dx : \varphi \in W_0^{1,p}(Q; \mathbb{R}^m) \right\}.$$

**8.1. Proof of Proposition 2.2.** The Proposition 2.2 follows from Proposition 8.1 by noticing that

$$\mathcal{Z}L(x, \xi) = \lim_{\varepsilon \rightarrow 0} \frac{m_\xi(Q_\varepsilon(x))}{\varepsilon^d},$$

and by using the following result.

**Lemma 8.6.** *If (2.4) holds then for a.a.  $x \in \Omega$  and every  $\xi \in \text{dom}L(x, \cdot)$  we have*

$$(8.7) \quad \lim_{\varepsilon \rightarrow 0} \frac{m_\xi(Q_\varepsilon(x))}{\varepsilon^d} \leq \lim_{\varepsilon \rightarrow 0} \frac{m_\xi^\#(Q_\varepsilon(x))}{\varepsilon^d}.$$

*Proof.* Fix  $\varepsilon \in ]0, 1[$  and  $s > 1$ . Fix  $x \in \Omega'$  where  $\Omega' = \{x \in \Omega : \text{dom}L(x, \cdot) \subset \Lambda_L(x)\}$  which satisfies  $|\Omega \setminus \Omega'| = 0$  since (2.4). Fix  $\xi \in \text{dom}L(x, \cdot)$  and fix  $\delta \in ]0, 2(s-1)\varepsilon^{1-d}[$ . Choose  $\{\overline{Q}_i\}_{i \geq 1} \in \mathcal{F}_{\frac{\delta\varepsilon^d}{2}}(Q_\varepsilon(x))$  such that  $|Q_\varepsilon(x) \setminus \bigcup_{i \geq 1} \overline{Q}_i| = 0$ ,  $Q_i \cap Q_\varepsilon(x) \neq \emptyset$  for all  $i \geq 1$ , and

$$(8.8) \quad \sum_{i \geq 1} m_\xi(Q_i) \leq \frac{\delta\varepsilon^d}{2} + m_\xi^\#(Q_\varepsilon(x)).$$

If  $O_\delta = \bigcup_{i \geq 1} Q_i$  then  $Q_\varepsilon(x) \subset O_\delta \subset Q_{s\varepsilon}(x)$ , indeed, on one hand we have  $O_\delta \subset \{y \in \Omega : \text{dist}(y, Q_\varepsilon(x)) < \frac{\delta\varepsilon^d}{2}\}$  and  $\frac{\delta\varepsilon^d}{2} + \varepsilon \leq s\varepsilon$ , thus  $O_\delta \subset Q_{s\varepsilon}(x)$ . On the other hand  $Q_\varepsilon(x) \setminus \overline{O}_\delta$  is open and  $|Q_\varepsilon(x) \setminus \overline{O}_\delta| \leq |Q_\varepsilon(x) \setminus O_\delta| = 0$ , therefore  $Q_\varepsilon(x) \setminus \overline{O}_\delta = \emptyset$ . It follows that  $Q_\varepsilon(x) \subset \overline{O}_\delta$  and so  $Q_\varepsilon(x) \subset O_\delta$ .



For each  $i \geq 1$  there exists  $\varphi_\delta^i \in W_0^{1,p}(\mathbb{Q}_i; \mathbb{R}^m)$  such that

$$(8.9) \quad \int_{\mathbb{Q}_i} L(y, \xi + \nabla \varphi_\delta^i) dy \leq \frac{\delta \varepsilon^d}{2^{i+1}} + m_\xi(\mathbb{Q}_i).$$

Define  $\varphi_\delta := \sum_{i \geq 1} \varphi_\delta^i \mathbb{I}_{\mathbb{Q}_i} \in W_0^{1,p}(O_\delta; \mathbb{R}^m)$ , then taking account of (8.8) we have

$$(8.10) \quad \begin{aligned} \int_{O_\delta} L(y, \xi + \nabla \varphi_\delta) dy &= \sum_{i \geq 1} \int_{\mathbb{Q}_i} L(y, \xi + \nabla \varphi_\delta^i) dy \leq \frac{\delta \varepsilon^d}{2} + \sum_{i \geq 1} m_\xi(\mathbb{Q}_i) \\ &\leq \delta \varepsilon^d + m_\xi^\sharp(\mathbb{Q}_\varepsilon(x)). \end{aligned}$$

The function  $\varphi_\delta$  also belongs in  $W_0^{1,p}(\mathbb{Q}_{s\varepsilon}(x); \mathbb{R}^m)$ , and moreover

$$(8.11) \quad \begin{aligned} \int_{O_\delta} L(y, \xi + \nabla \varphi_\delta) dy &\geq \int_{\mathbb{Q}_{s\varepsilon}(x)} L(y, \xi + \nabla \varphi_\delta) dy - \int_{\mathbb{Q}_{s\varepsilon}(x) \setminus \mathbb{Q}_\varepsilon(x)} L(y, \xi) dy \\ &\geq m_\xi(\mathbb{Q}_{s\varepsilon}(x)) - \int_{\mathbb{Q}_{s\varepsilon}(x) \setminus \mathbb{Q}_\varepsilon(x)} L(y, \xi) dy. \end{aligned}$$

From (2.4) we have

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^d} \int_{\mathbb{Q}_{s\varepsilon}(x) \setminus \mathbb{Q}_\varepsilon(x)} L(y, \xi) dy &= \overline{\lim}_{\varepsilon \rightarrow 0} \left\{ s^d \int_{\mathbb{Q}_{s\varepsilon}(x)} L(y, \xi) dy - \int_{\mathbb{Q}_\varepsilon(x)} L(y, \xi) dy \right\} \\ &\leq (s^d - 1)L(x, \xi). \end{aligned}$$

From (8.10) and (8.11) it holds since  $s > 1$

$$(8.12) \quad \overline{\lim}_{\varepsilon \rightarrow 0} \frac{m_\xi(\mathbb{Q}_{s\varepsilon}(x))}{(s\varepsilon)^d} \leq (s^d - 1)L(x, \xi) + \delta + \overline{\lim}_{\varepsilon \rightarrow 0} \frac{m_\xi^\sharp(\mathbb{Q}_\varepsilon(x))}{\varepsilon^d}.$$

But again since  $s > 1$  we have

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \frac{m_\xi(\mathbb{Q}_{s\varepsilon}(x))}{(s\varepsilon)^d} &= \lim_{\varepsilon \rightarrow 0} \sup_{\eta \in ]0, \varepsilon[} \frac{m_\xi(\mathbb{Q}_{s\eta}(x))}{(s\eta)^d} \\ &= \lim_{\varepsilon \rightarrow 0} \sup_{\eta' \in ]0, s\varepsilon[} \frac{m_\xi(\mathbb{Q}_{\eta'}(x))}{(\eta')^d} \\ &\geq \lim_{\varepsilon \rightarrow 0} \sup_{\eta' \in ]0, \varepsilon[} \frac{m_\xi(\mathbb{Q}_{\eta'}(x))}{(\eta')^d} = \overline{\lim}_{\varepsilon \rightarrow 0} \frac{m_\xi(\mathbb{Q}_\varepsilon(x))}{\varepsilon^d}. \end{aligned}$$

Therefore from (8.12) we obtain

$$\overline{\lim}_{\varepsilon \rightarrow 0} \frac{m_\xi(\mathbb{Q}_\varepsilon(x))}{\varepsilon^d} \leq (s^d - 1)L(x, \xi) + \delta + \overline{\lim}_{\varepsilon \rightarrow 0} \frac{m_\xi^\sharp(\mathbb{Q}_\varepsilon(x))}{\varepsilon^d},$$

send  $s \rightarrow 1$  and  $\delta \rightarrow 0$  we obtain (8.7) and the proof is complete.  $\blacksquare$

## REFERENCES

- [ABF03] E. Acerbi, G. Bouchitté, and I. Fonseca. Relaxation of convex functionals: the gap problem. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20(3):359–390, 2003.
- [AH96] David R. Adams and Lars Inge Hedberg. *Function spaces and potential theory*, volume 314 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1996.
- [AH10] Omar Anza Hafsa. On the integral representation of relaxed functionals with convex bounded constraints. *ESAIM Control Optim. Calc. Var.*, 16(1):37–57, 2010.
- [AHM07] Omar Anza Hafsa and Jean-Philippe Mandallena. Relaxation of variational problems in two-dimensional nonlinear elasticity. *Ann. Mat. Pura Appl. (4)*, 186(1):187–198, 2007.
- [AHM08] Omar Anza Hafsa and Jean-Philippe Mandallena. Relaxation theorems in nonlinear elasticity. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 25(1):135–148, 2008.

- [AHM11] Omar Anza Hafsa and Jean-Philippe Mandallena. Homogenization of nonconvex integrals with convex growth. *J. Math. Pures Appl. (9)*, 96(2):167–189, 2011.
- [AHM12] Omar Anza Hafsa and Jean-Philippe Mandallena. Homogenization of unbounded singular integrals in  $W^{1,\infty}$ . *to appear on Ric. Mat.*, 2012.
- [BB95] G. Buttazzo and M. Belloni. A survey on old and recent results about the gap phenomenon in the calculus of variations. In *Recent developments in well-posed variational problems*, volume 331 of *Math. Appl.*, pages 1–27. Kluwer Acad. Publ., Dordrecht, 1995.
- [BB00a] Hamed Ben Belgacem. Relaxation of singular functionals defined on Sobolev spaces. *ESAIM Control Optim. Calc. Var.*, 5:71–85 (electronic), 2000.
- [BB00b] Guy Bouchitté and Michel Bellieud. Regularization of a set function—application to integral representation. *Ricerche Mat.*, 49(suppl.):79–93, 2000. Contributions in honor of the memory of Ennio De Giorgi (Italian).
- [BD98] Andrea Braides and Anneliese Defranceschi. *Homogenization of multiple integrals*, volume 12 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press Oxford University Press, New York, 1998.
- [BFM98] Guy Bouchitté, Irene Fonseca, and Luisa Mascarenhas. A global method for relaxation. *Arch. Rational Mech. Anal.*, 145(1):51–98, 1998.
- [BM84] J. M. Ball and F. Murat.  $W^{1,p}$ -quasiconvexity and variational problems for multiple integrals. *J. Funct. Anal.*, 58(3):225–253, 1984.
- [Bon72] Benedetto Bongiorno. Sulla derivabilità delle funzioni arbitrarie d’insieme. *Rend. Circ. Mat. Palermo (2)*, 21:71–84, 1972.
- [But89] Giuseppe Buttazzo. *Semicontinuity, relaxation and integral representation in the calculus of variations*, volume 207 of *Pitman Research Notes in Mathematics Series*. Longman Scientific & Technical, Harlow, 1989.
- [CDA02] Luciano Carbone and Riccardo De Arcangelis. *Unbounded functionals in the calculus of variations*, volume 125 of *Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics*. Chapman & Hall/CRC, Boca Raton, FL, 2002. Representation, relaxation, and homogenization.
- [Dac08] Bernard Dacorogna. *Direct methods in the calculus of variations*, volume 78 of *Applied Mathematical Sciences*. Springer, New York, second edition, 2008.
- [DAMZ04] Riccardo De Arcangelis, Sara Monsurrò, and Elvira Zappale. On the relaxation and the Lavrentieff phenomenon for variational integrals with pointwise measurable gradient constraints. *Calc. Var. Partial Differential Equations*, 21(4):357–400, 2004.
- [DAZ05] Riccardo De Arcangelis and Elvira Zappale. The relaxation of some classes of variational integrals with pointwise continuous-type gradient constraints. *Appl. Math. Optim.*, 51(3):251–277, 2005.
- [DGS73] E. De Giorgi and S. Spagnolo. Sulla convergenza degli integrali dell’energia per operatori ellittici del secondo ordine. *Boll. Un. Mat. Ital. (4)*, 8:391–411, 1973.
- [DMM86] Gianni Dal Maso and Luciano Modica. Integral functionals determined by their minima. *Rend. Sem. Mat. Univ. Padova*, 76:255–267, 1986.
- [Fed69] Herbert Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer-Verlag New York Inc., New York, 1969.
- [FM92] Irene Fonseca and Stefan Müller. Quasi-convex integrands and lower semicontinuity in  $L^1$ . *SIAM J. Math. Anal.*, 23(5):1081–1098, 1992.
- [HK60] W. E. Hartnett and A. H. Kruse. Differentiation of set functions using Vitali coverings. *Trans. Amer. Math. Soc.*, 96:185–209, 1960.
- [Mat95] Pertti Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability.
- [Zap05] Elvira Zappale. Relaxation in presence of pointwise gradient constraints. *Ricerche Mat.*, 54(2):655–660 (2006), 2005.

UNIVERSITE DE NIMES, LABORATOIRE MIPA, SITE DES CARMES, PLACE GABRIEL PÉRI,  
30021 NÎMES, FRANCE

LABORATOIRE LMGC, UMR-CNRS 5508, PLACE EUGÈNE BATAILLON, 34095 MONTPELLIER,  
FRANCE.

*E-mail address:* Omar Anza Hafsa <omar.anza-hafsa@unimes.fr>

UNIVERSITE DE NIMES, LABORATOIRE MIPA, SITE DES CARMES, PLACE GABRIEL PÉRI,  
30021 NÎMES, FRANCE

*E-mail address:* Jean-Philippe Mandallena <jean-philippe.mandallena@unimes.fr>